

FULL PAPER

Domination topological properties of some chemical structures using φ_p -Polynomial approach

Hanan Ahmed^a  | Mohammad Reza Farahani^{b,*}  | Anwar Alwardi^c  | Ruby Salestina M.^a

^aDepartment of Mathematics, Yuvaraja's college, University of Mysore, Mysuru, India

^bDepartment of Applied Mathematics, Iran University of Science and Technology (IUST), Narmak, Tehran 16846, Iran

^cDepartment of Mathematics, University of Aden, Yemen

φ_p -Polynomial is one way to represent a graph algebraically, and it has a major role in theoretical chemistry. It is used in the calculation of the exact values of many topological indices that depend on the P set degree. In this paper, we study the φ_p -Polynomial by using minimal and minimum dominating sets for Nicotine, Aspirin, and Anthraquinone. Using those φ_p -Polynomials, some domination and domination topological indices are derived. Also, the results are graphically interpreted.

***Corresponding Author:**

Mohammad Reza Farahani
Email: mrfarahani88@gmail.com
Tel.: +989192478265

KEYWORDS

φ_p -Polynomial; domination topological indices; anthraquinone; nicotine; aspirin.

Introduction

Chemical graph theory is one of the branches of mathematical chemistry, where it is important and necessary for a better understanding and interpretation of the nature of the chemical composition. Recently, the chemical graph theory is used in organizing and arranging the existing problem, because it provides an arrangement of the rules and laws according to a specific systems and specific planning. Topological indices can be viewed as molecular descriptors that describe the composition of chemical structures and help predict some of the chemical and physical properties of these structures. Anthraquinone is an aromatic organic compound with the formula $C_{14}H_8O_2$. It is a highly crystalline yellow solid that is poorly soluble in water but soluble in hot organic solvents. Hydrogenation gives dihydroanthraquinone(anthrahydroquinone) [13]. Sulfonation with sulfuric acid gives anthraquinone-1-sulfonic acid [16], which

reacts with sodium chlorate to give 1-chloro-anthaquinone [17]. Nicotine is a widely used stimulant and a parasympathetic alkaloid, as it is produced naturally from the roots of the nicotine talcum plant, a plant found in the American continent, where it was extracted from the leaves of this plant in 1828 [18, 19, 22]. The pure state of nicotine is an alkaloid that has no color, is volatile, soluble in water, and is also considered bioactive [22] it can also be absorbed through the skin and mucous membranes in the mouth and nose, as well as in the alveoli [2]. The largest nicotine consumption can be considered through cigarettes, which contain an average of 14 mg per cigarette. It is noted that the amount provided by smoking varies and depends on the quality of the cigarettes and other factors [20]. The cerebral half-life of nicotine is 52 minutes [1, 14, 19, 22, 24-35]. Aspirin is considered one of the oldest medicines due to its frequent use in various fields of medicine. Aspirin was accepted for back pain in 1903 and in 1923 aspirin was

used to treat headaches, and it was also used for arthritis in 1933. Studies in 1988 showed that aspirin could be effective in preventing and treating gallstones [10]. Also, there are many studies that have shown beneficial effects on various cancer cases see [3, 4, 11, 15, 21]. Aspirin is considered an anti-headache and insomnia, a pain reliever, and an anti-fever in the case of infectious diseases and is against blood clots. A set $D \subseteq V$ is said to be a dominating set of G , if for any vertex $v \in V - D$ there exists a vertex $u \in D$ such that u and v are adjacent. A dominating-set $D = \{v_1, v_2, \dots, v_r\}$ is minimal if $D - v_i$ is not a dominating set [12]. A dominating set of G of minimum cardinality is said to be a minimum dominating set. A topological index is a numerical parameter of the graph, such that this parameter is the same for the graph which is isomorphism.

Basic definitions

TABLE 1 Description of some domination and γ -domination topological indices

D indices	$f(d_d(u), d_d(v))$	γ D indices	$f(d_\gamma(u), d_\gamma(v))$
$DM_1^*(G)$	$d_d(u) + d_d(v)$	$\gamma M_1^*(G)$	$d_\gamma(u) + d_\gamma(v)$
$DF^*(G)$	$d_d^2(u) + d_d^2(v)$	$\gamma F^*(G)$	$d_\gamma^2(u) + d_\gamma^2(v)$
$DM_2(G)$	$d_d(u) + d_d(v)$	$\gamma M_2(G)$	$d_\gamma(u)d_\gamma(v)$
$HD(G)$	$d_d^2(u) + d_d^2(v) + 2d_d(u)d_d(v)$	$\gamma H(G)$	$d_\gamma^2(u) + d_\gamma^2(v) + 2d_\gamma(u)d_\gamma(v)$

TABLE 2 Derivation of domination and domination topological indices from ϕ_P -Polynomials

D indices	Derivation from $\phi_d(G)$	γ D indices	Derivation from $\phi_d(G)$
$DM_1^*(G)$	$(D_x + D_y)(\phi_d(G)) _{x=y=1}$	$\gamma M_1^*(G)$	$(D_x + D_y)(\phi_\gamma(G)) _{x=y=1}$
$DF^*(G)$	$(D_x^2 + D_y^2)(\phi_d(G)) _{x=y=1}$	$\gamma F^*(G)$	$(D_x^2 + D_y^2)(\phi_\gamma(G)) _{x=y=1}$
$DM_2(G)$	$(D_x D_y)(\phi_d(G)) _{x=y=1}$	$\gamma M_2(G)$	$(D_x D_y)(\phi_\gamma(G)) _{x=y=1}$
$HD(G)$	$(D_x^2 + D_y^2 + 2D_x D_y)(\phi_d(G)) _{x=y=1}$	$\gamma H(G)$	$(D_x^2 + D_y^2 + 2D_x D_y)(\phi_\gamma(G)) _{x=y=1}$

$$\text{Here, } D_x(f(x, y)) = x \frac{\partial(f(x, y))}{\partial x},$$

$$D_y(f(x, y)) = y \frac{\partial(f(x, y))}{\partial y}.$$

In [7-9] authors have introduced new topological indices called domination and γ -

In Hanan Ahmed et al. [6] the definition of ϕ_P -polynomial is introduced as:

Definition 2.1. Let $G=(V,E)$ be a graph, $d_P(v)$ be the P set degree of the vertex v denoted by $d_P(v)=|\{S \subseteq V(G): S \text{ has property } P \text{ and } v \in S\}|$.

The minimum and maximum P set degree of G denote as $\delta_P(G)=\delta_P$ and $\Delta_P(G)=\Delta_P$ respectively. Such that $\delta_P=\min\{d_P(v): v \in V(G)\}$ and $\Delta_P=\max\{d_P(v): v \in V(G)\}$.

Let $d_{Pm_{i,j}}(G)=|\{e=uv: d_P(u)=i, d_P(v)=j\}|$. The ϕ_P -polynomial is defined as

$$\phi_P(G, x, y) = \sum_{\delta_P \leq i \leq j \leq \Delta_P} d_{Pm_{i,j}}(G) x^i y^j.$$

Domination (D) and γ -Domination (γ D) indices defined on $E(G)$, which can be written as TABLE 1 follows:

$$D(G) = \sum_{uv \in E(G)} f(d_d(u), d_d(v)),$$

$$\gamma D(G) = \sum_{uv \in E(G)} f(d_\gamma(u), d_\gamma(v)).$$

domination topological indices, which are defined as

$$\begin{aligned} DM_1(G) &= \sum_{v \in V(G)} d_d^2(v), \\ DM_2(G) &= \sum_{uv \in E(G)} d_d(u)d_d(v), \\ DM_1^* &= \sum_{uv \in E(G)} (d_d(u) + d_d(v)), \\ DF^*(G) &= \sum_{uv \in E(G)} d_d^2(u) + d_d^2(v), \\ DF(G) &= \sum_{v \in V(G)} d_d^3(v), \\ DH(G) &= \sum_{uv \in E(G)} (d_d(u) + d_d(v))^2. \end{aligned}$$

Where $d_d(v)$ is the domination degree of $v \in V(G)$ and defined as the number of minimal dominating sets of G which contains v . The minimum and maximum domination degree of G are denoted by $\delta_d(G)=\delta_d$ and $\Delta_d(G)=\Delta_d$, respectively, in which $\delta_d=\min\{d_d(v): v \in V(G)\}$ and $\Delta_d=\max\{d_d(v): v \in V(G)\}$.

Definition 2.2. [5] For all vertex $v \in V(G)$, the domination value of v define as: $d_\gamma(v)=|\{S \subseteq V(G): S \text{ is a minimum dominating set and } v \in S\}|$.

We denote the minimum and maximum domination value of a graph G by: $\delta_\gamma(G)=\delta_\gamma=\min\{d_\gamma(v): v \in V(G)\}$ and $\Delta_\gamma(G)=\Delta_\gamma=\max\{d_\gamma(v): v \in V(G)\}$ respectively.

The γ -domination Zagreb, γ -domination forgotten, γ -domination hyper indices are defined as:

$$\begin{aligned} \gamma M_1(G) &= \sum_{v \in V(G)} d_\gamma^2(v), \\ \gamma M_2(G) &= \sum_{uv \in E(G)} d_\gamma(u)d_\gamma(v), \\ \gamma F(G) &= \sum_{v \in V(G)} d_\gamma^3(v), \\ \gamma H(G) &= \sum_{uv \in E(G)} (d_\gamma(u) + d_\gamma(v))^2, \\ \gamma M_1^* &= \sum_{uv \in E(G)} d_\gamma(u) + d_\gamma(v), \\ \gamma F^*(G) &= \sum_{uv \in E(G)} d_\gamma^2(u) + d_\gamma^2(v). \end{aligned}$$

Results and discussion: Anthraquinone

TABLE 3 Domination degree of the vertices of G . (*)

$d_d(v)$	50	40	30
Number of the vertices	4	9	3

(*) To obtain φd and $\varphi \gamma$ polynomials of $G \approx$ anthraquinone, Tables 3 and 4 are very essential

TABLE 4 Domination value of the vertices of G

$d_\gamma(v)$	35	32	25	22	21	16	14
Number of the vertices	1	1	3	2	3	4	2

Lemma 3.1. The total number of minimal and minimum dominating sets in the molecular graph of anthraquinone is 100 and 57, respectively.




FIGURE1 Anthraquinone

Proof. Let G be the molecular graph of anthraquinone. We first divide G into three components: C_1 , C_2 , and C_3 . We calculate the minimal dominating sets of each component, so that we get $T_m(C_1)=5$, $T_m(C_2)=7$, and $T_m(C_3)=5$.

Every minimal dominating set of C_1 is added to each minimal dominating set of C_2 , and we check for the minimality of the resulting dominating sets. As a result, we get $5 \times 7 = 35$ minimal dominating sets, of which 10 are repeated. Hence, we get 25 minimal dominating sets.

Again, every minimal dominating set of C_3 is added to each of the 25 minimal dominating sets, and we check for the minimality of the resulting dominating sets. In this case, we get a total of $5 \times 25 = 125$ minimal dominating sets, of which 25 are repeated. In all, there are 100 minimal dominating sets of G . Note that $\gamma(G)=6$, and we have the fact that every minimum dominating set is a minimal dominating set. Out of 100 minimal dominating sets, there are exactly 57 sets whose cardinality is equal to the domination number of G . Hence, G has 57 minimum dominating sets. ■

Theorem 3.2. If G is the molecular graph of anthraquinone, then

$$\varphi_d(G, x, y) = x^{30}[y^{30} + 4y^{40} + 3y^{50}] + x^{40}[7y^{40} + y^{50}] + 2x^{50}y^{50}.$$

$$\varphi_\gamma(G, x, y) = x^{14}[y^{14} + y^{21} + y^{22} + y^{32} + y^{35}] + x^{16}[2y^{16} + 4y^{25} + y^{32} + y^{35}] + x^{21}[3y^{22} + y^{35}] + x^{25}y^{32}.$$

TABLE 5 Edges partition

$d_{dm_{ij}}(G)$	(30,30)	(30,40)	(30,50)	(40,40)	(40,50)	(50,50)
Number of the edges	1	4	3	7	1	2

Hence from Table 5, we get

$$\begin{aligned}\varphi_d(G, x, y) &= \sum_{\delta_d \leq i \leq j \leq \Delta_d} d_{dm_{ij}}(G) x^i y^j \\ &= x^{30}[y^{30} + 4y^{40} + 3y^{50}] + x^{40}[7y^{40} + y^{50}] + 2x^{50}y^{50}.\end{aligned}$$

TABLE 6 Edge partition

Number of edges	$d_\gamma m_{ij}$
1	(14,14), (14,21), (14,22), (14,32), (14,35), (16,32), (16,35), (21,35), (25,32)
2	(16,16)
3	(21,22)
4	(16,25)

So, from Table 6, we get

$$\varphi_\gamma(G, x, y) = \sum_{\delta_\gamma \leq i \leq j \leq \Delta_\gamma} d_\gamma m_{ij}(G) x^i y^j = x^{14}[y^{14} + y^{21} + y^{22} + y^{32} + y^{35}] + x^{16}[2y^{16} + 4y^{25} + y^{32} + y^{35}] + x^{21}[3y^{22} + y^{35}] + x^{25}y^{32} \blacksquare$$





FIGURE 2 Plotting of (a) φ_d -polynomial and (b) φ_γ -polynomial of Anthraquinone

Theorem 3.3. Suppose G is the molecular graph of anthraquinone, then

1. $DM_1^*(G) = 1430, \gamma M_1(G) = 763,$
2. $DF^*(G) = 58500, \gamma F^*(G) = 17749,$
3. $DM_2(G) = 28400, \gamma M_2(G) = 7841,$
4. $DH(G) = 115300, \gamma H(G) = 33431.$

Proof. We have

$$\begin{aligned}\varphi_d(G, x, y) &= x^{30}[y^{30} + 4y^{40} + 3y^{50}] + x^{40}[7y^{40} + y^{50}] \\ &\quad + 2x^{50}y^{50}.\end{aligned}$$

Then

$$\begin{aligned}(D_x + D_y)(\varphi_d(G, x, y)) &= x^{30}[60y^{30} + 280y^{40} + 240y^{50}] \\ &\quad + x^{40}[560y^{40} + 90y^{50}] + 200x^{50}y^{50},\end{aligned}$$

$$\begin{aligned}(D_x^2 + D_y^2)(\varphi_d(G, x, y)) &= x^{30}[1800y^{30} + 10000y^{40} + 10200y^{50}] + x^{40}[22400y^{40} + 4100y^{50}] \\ &\quad + 10000x^{50}y^{50}, \\ (D_x D_y)(\varphi_d(G, x, y)) &= x^{30}[900y^{30} + 4800y^{40} + 4500y^{50}] \\ &\quad + x^{40}[11200y^{40} + 2000y^{50}] + 5000x^{50}y^{50}, \\ (D_x + D_y)^2(\varphi_d(G, x, y)) &= x^{30}[3600y^{30} + 19600y^{40} + 19200y^{50}] \\ &\quad + x^{40}[44800y^{40} + 8100y^{50}] + 20000x^{50}y^{50}\end{aligned}$$

By using Table 2, we get

$$\begin{aligned}DM_1^*(G) &= x^{30}[60y^{30} + 280y^{40} + 240y^{50}] + x^{40}[560y^{40} + 90y^{50}] + 200x^{50}y^{50}|_{x=y=1} = 1430, \\ DF^*(G) &= x^{30}[1800y^{30} + 10000y^{40} + 10200y^{50}] + x^{40}[22400y^{40} + 4100y^{50}] + 10000x^{50}y^{50}|_{x=y=1} =\end{aligned}$$

58500,

$$\begin{aligned} DM_2(G) = & x^{30}[900y^{30} + 4800y^{40} + 4500y^{50}] \\ & + x^{40}[11200y^{40} + 2000y^{50}] + 5000x^{50}y^{50}|_{x=y=1} = \\ & 28400, \end{aligned}$$

$$\begin{aligned} DH(G) = & x^{30}[3600y^{30} + 19600y^{40} + 19200y^{50}] \\ & + x^{40}[44800y^{40} + 8100y^{50}] + 20000x^{50}y^{50}|_{x=y=1} = \\ & 115300; \end{aligned}$$

For γ -domination indices we have,

$$\begin{aligned} \varphi_\gamma(G, x, y) = & x^{14}[y^{14} + y^{21} + y^{22} + y^{32} + y^{35}] \\ & + x^{16}[2y^{16} + 4y^{25} + y^{32} + y^{35}] + x^{21}[3y^{22} + y^{35}] + x^{25} + y^{32} \end{aligned}$$

Then

$$\begin{aligned} (D_x + D_y)(\varphi_\gamma(G, x, y)) = & x^{14}[28y^{14} + 35y^{21} + 36y^{22} \\ & + 46y^{32} + 49y^{35}] + x^{16}[64y^{16} + 164y^{25} + 48y^{32} + 51y^{35}] \\ & + x^{21}[129y^{22} + 56y^{35}] + 57x^{25}y^{32}, \end{aligned}$$

$$\begin{aligned} (D_x^2 + D_y^2)(\varphi_\gamma(G, x, y)) = & x^{14}[392y^{14} + 637y^{21} + 680y^{22} \\ & + 1220y^{32} + 1421y^{35}] + x^{16}[1024y^{16} + 3524y^{25} \\ & + 1280y^{32} + 1481y^{35}] + x^{21}[2775y^{22} + 1666y^{35}] \\ & + 1649x^{25}y^{32}, \end{aligned}$$

$$\begin{aligned} (D_x D_y)(\varphi_\gamma(G, x, y)) = & x^{14}[196y^{14} + 294y^{21} + 308y^{22} \\ & + 448y^{32} + 490y^{35}] + x^{16}[512y^{16} + 1600y^{25} \\ & + 512y^{32} + 560y^{35}] + x^{21}[1386y^{22} + 735y^{35}] + 800x^{25} \\ & y^{32}, \end{aligned}$$

$$\begin{aligned} (D_x^2 + D_y^2 + D_x D_y)(\varphi_\gamma(G, x, y)) = & x^{14}[784y^{14} + 1225y^{21} \\ & + 1296y^{22} + 2116y^{32} + 2401y^{35}] + x^{16}[2048y^{16} \\ & + 6724y^{25} + 2304y^{32} + 2601y^{35}] + x^{21}[5547y^{22} \\ & + 3136y^{35}] + 3249x^{25}y^{32} \end{aligned}$$

By using Table 2, we get

$$\begin{aligned} M_1^*(G) = & x^{14}[28y^{14} + 35y^{21} + 36y^{22} + 46y^{32} + 49y^{35}] \\ & + x^{16}[64y^{16} + 164y^{25} + 48y^{32} + 51y^{35}] \\ & + x^{21}[129y^{22} + 56y^{35}] + 57x^{25}y^{32}|_{x=y=1} = 763, \end{aligned}$$

$$\begin{aligned} F^*(G) = & x^{14}[392y^{14} + 637y^{21} + 680y^{22} + 1220y^{32} \\ & + 1421y^{35}] + x^{16}[1024y^{16} + 3524y^{25} + 1280y^{32} \\ & + 1481y^{35}] + x^{21}[2775y^{22} + 1666y^{35}] \\ & + 1649x^{25}y^{32}|_{x=y=1} = 17749, \end{aligned}$$

$$\begin{aligned} M_2(G) = & x^{14}[196y^{14} + 294y^{21} + 308y^{22} + 448y^{32} \\ & + 490y^{35}] + x^{16}[512y^{16} + 1600y^{25} + 512y^{32} + 560y^{35}] \\ & + x^{21}[1386y^{22} + 735y^{35}] + 800x^{25}y^{32}|_{x=y=1} = 7841, \end{aligned}$$

$$\begin{aligned} H(G) = & x^{14}[784y^{14} + 1225y^{21} + 1296y^{22} + 2116y^{32} \\ & + 2401y^{35}] + x^{16}[2048y^{16} + 6724y^{25} + 2304y^{32} + 2601 \\ & y^{35}] + x^{21}[5547y^{22} + 3136y^{35}] + 3249x^{25}y^{32}|_{x=y=1} \end{aligned}$$

$$= 33431.$$

Results and discussion: Nicotin

Lemma 4.1. Suppose G is the molecular graph of the nicotine. The total number of a

minimal and minimum dominating set is 29 and 6, respectively. To obtain φ_d and φ_γ -polynomials of G, Tables 7 and 8 are very essential.

TABLE 7 Domination degree of vertices of G.

$d_d(v)$	9	10	11	12	14	19
Number of vertices	1	3	2	3	2	1

TABLE 8 Domination value of vertices of G

$d_\gamma(v)$	0	2	3	6
Number of vertices	3	6	2	1




FIGURE 3 Nicotin

Theorem 4.2. Let G be the molecular graph of nicotine. Then:

(i)

$$\varphi_d(G, x, y) = x^9[2y^{10} + y^{14}] + x^{10}[y^{10} + 2y^{12} + y^{14} + y^{19}] + 4x^{11}y^{12} + x^{14}y^{14},$$

$$(ii) \varphi_\gamma(G, x, y) = y^2 + 2y^3 + 3y^6 + 6x^2y^2 + x^3y^3$$

Proof. Case 1. The set of edges that is ended by domination degree can be divided into:

$$d_dm_{9,10}=2, \quad d_dm_{9,14}=1, \quad d_dm_{10,10}=1, \quad d_dm_{10,12}=2, \\ d_dm_{10,14}=1, \quad d_dm_{10,19}=1, \quad d_dm_{11,12}=4, \quad d_dm_{14,14}=1.$$

Now

$$\begin{aligned} \varphi_d(G, x, y) = & \sum_{\delta_d \leq i \leq j \leq \Delta_d} d_dm_{i,j}(G)x^i y^j \\ = & d_dm_{9,10}x^9y^{10} + d_dm_{9,14}x^9y^{14} + d_dm_{10,10}x^{10}y^{10} + \\ & d_dm_{10,12}x^{10}y^{12} + d_dm_{10,14}x^{10}y^{14} + d_dm_{10,19}x^{10}y^{19} \\ & + d_dm_{11,12}x^{11}y^{12} + d_dm_{14,14}x^{14}y^{14} \end{aligned}$$

After putting the value of $d_dm_{i,j}$, we get

$$\begin{aligned} \varphi_d(G, x, y) = & x^9[2y^{10} + y^{14}] + x^{10}[y^{10} + 2y^{12} + y^{14} + y^{19}] + \\ & 4x^{11}y^{12} + x^{14}y^{14} \end{aligned}$$

Case 2: Likewise, the edges whose beginning and end are domination value can be divided as:

$$d_\gamma m_{0,2}=1, d_\gamma m_{0,3}=2, d_\gamma m_{0,6}=3, d_\gamma m_{2,2}=6, d_\gamma m_{3,3}=1.$$

As above we get

$$\varphi_Y(G,x,y)=y^2+2y^3+3y^6+6x^2y^2+x^3y^3. \blacksquare$$

Theorem 4.3. Suppose G is the molecular graph of nicotine. Then, we have

1. $DM_1^*(G)=298, \gamma M_1(G)=59,$
2. $DF^*(G)=3536, \gamma F^*(G)=196,$
3. $DM_2(G)=1700, \gamma M_2(G)=33,$
4. $DH(G)=6936, \gamma H(G)=262:$




FIGURE 4 Plotting of (a) φ_d -polynomial and (b) φ_Y -polynomial of Nicotine

Proof. We have

$$\begin{aligned} \varphi_d(G,x,y) &= x^9[2y^{10}+y^{14}]+x^{10}[y^{10}+2y^{12}+y^{14}+y^{19}] \\ &+ 4x^{11}y^{12}+x^{14}y^{14}, \end{aligned}$$

$$\varphi_Y(G,x,y)=y^2+2y^3+3y^6+6x^2y^2+x^3y^3;$$

Then

$$\begin{aligned} (D_x+D_y)(\varphi_d(G,x,y)) &= x^9[38y^{10}+23y^{14}]+x^{10}[20y^{10} \\ &+ 44y^{12}+24y^{14}+29y^{19}]+92x^{11}y^{12}+28x^{14}y^{14}, \end{aligned}$$

$$(D_x^2+D_y^2)(\varphi_d(G,x,y))=x^9[362y^{10}+277y^{14}]$$

$$\begin{aligned} &+x^{10}[200y^{10}+488y^{12}+296y^{14}+461y^{19}] \\ &+1060x^{11}y^{12}+392x^{14}y^{14}, \end{aligned}$$

$$\begin{aligned} (D_xD_y)(\varphi_d(G,x,y)) &= 9x^9[20y^{10}+14y^{14}] \\ &+10x^{10}[10y^{10}+24y^{12}+14y^{14}+19y^{19}] \end{aligned}$$

$$+528x^{11}y^{12}+196x^{14}y^{14},$$

$$\begin{aligned} (D_x^2+D_y^2+2D_xD_y)(\varphi_d(G,x,y)) &= x^9[722y^{10}+529y^{14}] \\ &+x^{10}[400y^{10}+968y^{12}+576y^{14}+841y^{19}] \\ &+2116x^{11}y^{12}+784x^{14}y^{14}, \\ (D_x+D_y)(\varphi_Y(G,x,y)) &= 2y^2+6y^3+18y^6+24x^2y^2+9x^3y^3, \end{aligned}$$

$$\begin{aligned} (D_x^2+D_y^2)(\varphi_Y(G,x,y)) &+ 4y^2+18y^3+108y^6 \\ &+ 48x^2y^2+18x^3y^3, \\ (D_xD_y)(\varphi_Y(G,x,y)) &= 24x^2y^2+9x^3y^3, \\ (D_x^2+D_y^2+2D_xD_y)(\varphi_Y(G,x,y)) &= 4y^2+18y^3 \\ &+ 108y^6+69x^2y^2+36x^3y^3. \end{aligned}$$

The required result can be obtained using Table 2.■

Results and discussion: Asprini

Lemma 5.1. Let G be the molecular graph of Aspirin. Then the total number of minimal and minimum dominating sets is 20 and 4, respectively.




FIGURE 5 Asprini

To obtain φ_d and φ_Y polynomials of G , molecular graph of Aspirin, Tables 9 and 10 are very essential.

TABLE 9 Domination degree of the vertices of G

$d_d(v)$	6	7	8	10
Number of the vertices	1	1	4	7

TABLE 10 Domination value of the vertices of G

$d_Y(v)$	0	1	2	4
Number of the vertices	6	3	2	2

Theorem 5.2. If G is the molecular graph of Aspirin, then

$$(i) \varphi_d(G,x,y)=x^6[y^7+y^8+y^{10}]+x^7[y^8+y^{10}]+3x^8y^8$$

$+5x^{10}y^{10}$,
 (ii) $\varphi_Y(G,x,y)=y^9[1+x+x^2]+y^4[5+x]+xy+3xy^2$.

Proof. The edges of the graph can be divided by depending on the domination degree and

TABLE 11 Edges partition

Number of edges	$d_{dm_{ij}}(i,j)$
1	6,7 & 6,8 & 6,10 & 7,8 & 7,10 & 0,9 & 1,1 & 1,4 & 1,9 & 2,9
3	8,8 & 1,2
5	10,10 & 0,4

We get the desired result through definition of φ_P -polynomial and Table 11.■




FIGURE 6 Plotting of (a) φ_d -polynomial and (b) φ_Y -Polynomial of Aspirin

Theorem 5.3. Let G be the molecular graph of Aspirin, then

1. $DM_1^*(G)=223$, $M_1^*(G)=66$,
2. $DF^*(G)=1967$, $F^*(G)=359$,
3. $DM_2(G)=968$, $M_2(G)=38$,
4. $DH(G)=3903$, $H(G)=438$.

Proof. We have

- (i) $\varphi_d(G,x,y)=x^6[y^7+y^8+y^{10}]+x^7[y^8+y^{10}]$
 $+3x^8y^8+5x^{10}y^{10}$,
 (ii) $\varphi_Y(G,x,y)=y^9[1+x+x^2]+y^4[5+x]+xy+3xy^2$.

By applying the same method of proof of Theorem 4.3, we get the results.■

the domination value of the ends, as represented in the Table 11 below.

Conclusion

We have studied and computed the properties of some chemical compounds namely anthraquinone, nicotine, and aspirin through domination and gamma domination topological indices. First, we found φ_d polynomial and φ_Y polynomial and their respective 3D graphs (Figures 2, 4 and 6). Then we computed the domination and gamma domination indices from these polynomials. It is known that topological indices have the ability to predict some different characteristics, such as the central factor, stability of chemical compounds, boiling point, etc.

Acknowledgements

The authors would like to thank the reviewers for their helpful suggestions and comments.

Orcid:

Hanán Ahmed: <https://orcid.org/0000-0002-4008-4873>

Mohammad Reza Farahani:
<https://orcid.org/0000-0003-2969-4280>
 Anwar Alwardi: <https://orcid.org/0000-0002-1908-6006>

References

- [1] T. Alkam, T. Nabeshima, *Neuroch. Int.*, **2019**, 125, 117-126. PMID: 30779928. [crossref], [Google Scholar], [Publisher]
- [2] L.K. Baraona, D. Lovelace, J.L. L. Daniels, Mc Daniel, *J. Midwifery. Womens Health*, **2017**,

- 62, 253-269. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [3] A.J.E. Bardia, C.M. Olson, D. Vachon, R.A. Lazovich, A.H. Vierkant., P.J. Wang, K.E. Limburg, J.R. Anderson, *Cancer. Breast. Cancer. Res. Treat.*, **2011**, *126*, 149-55. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [4] P.K. Dhillon, S.A. Kenfield, M.J. Stampfer, E.L. Giovannuccilnt, *J. Cancer.*, **2011**, *128*, 2444-2452. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [5] Y. Eunjeong, arXiv,1109.6277v2 [math.CO] 1 Mar **2012**.
- [6] H. Ahmed, A. Alwardi, M. R. Salestina M., S. Nandappa D., *Biointerface Res. Appl. Chem.*, **2021**, *11*, 13290-13302. arXiv,1109.6277v2 [math.CO] 1 Mar **2012**. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [7] H. Ahmed, A. Alwardi, R. Salestina M., *Int.l J. Anal. Appl.*, **2020**, *19*, 47-64. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [8] H. Ahmed, A. Alwardi, R. Salestina M., N.D. Soner, *J. Dis. Math. Sci. Cryp.*, **2021**, *24*, 325-341.
- [9] A. Hanan, A. Alwardi, R. Salestina M., *J. Dis. Math. Sci. Cryp.*, **2021**, *24*, 353-368.
- [10] F.R. Heller, *Acta Gastroenterol Belg.*, **1990**, *53*, 16-21. PMID: 2114725.
- [11] L. Mascitelli, F. Pezzetta, J.L. Sullivan, *Med Hypotheses*, **2010**, *74*, 78-80. PMID: 19692186. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [12] J.F. Couturier, R. Letourneur, M. Liedlo, *Theoretica Com. Sci.*, **2015**, *562*, 634-642. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [13] L.C. Macleod, C.F. Allen, H. Benzanthrone, *Org. Synth.*, **1934**, *14*, 4-5. <https://doi.org/10.15227/orgsyn.014.0004>
- [14] A. Marsot, N. Simon, *Int. J. Toxicol.*, **2016**, *35*, 179-185. PMID: 26681385. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [15] P.M. Rothwell, F.G. Fowkes, J.F. Belch, H. Ogawa, C.P. Warlow, T.W. Meade, *Lancet.*, **2011**, *377*, 31-41. PMID: 21144578. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [16] W.J. Scott, C.F.H. Allen, *Org. Synth.*, **1938**, *18*, 72-73. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [17] W.J. Scott, C.F.H. Allen, *Org. Synth.*, **1938**, *18*, 15-16. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [18] M. Sieber, *Curr Med Chem.*, **2012**, *19*, 292-297. PMID: 22320303. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [19] L.M. Siqueira, *Pediatrics.*, **2017**, *139*, 1-15. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [20] G. Valentine, M. Sofuooglu, *Cur Neuropharmacol.*, **2018**, *16*, 403-414. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [21] A.L. Van Dyke, M.L. Cote, G. Prysak, G.B. Claeys, A.S. Wenzlaff, A.G. Schwartz., *Cancer Epid. Biomark. Prev.*, **2008**, *17*, 148-57. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [22] M. Zarrindasta, F. Khakpaei, *Brain. Research.*, **2019**, *1710*, 92-101.
- [23] D. Afzal, S. Hussain, M. Aldemir, M. Farahani, F. Afzal, *Eurasian Chem. Commun.*, **2020**, *2*, 1117-1125. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [24] M. Cancan, S. Ediz, M.R. Farahani, *Eurasian Chem. Commun.*, **2020**, *2*, 641-645. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [25] M. Alaeiyan, C. Natarajan, G. Sathiamoorthy, M.R. Farahani, *Eurasian Chem. Commun.*, **2020**, *2*, 646-651. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [26] M.R. Farahani, *Acta Chim. Slov.*, **2012**, *59*, 965-968. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- <https://pubmed.ncbi.nlm.nih.gov/24061384>
- [27] Z. Shao, A.R. Virk, M.S. Javed, M.A. Rehman, M.R. Farahani, *Eng. Appl. Sci. Lett.*, **2019**, *2*, 01-11. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [28] M.R. Farahani, *World Appl. Sci. J.*, **2012**, *20*, 1248-1251. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [29] H. Siddiqui, M.R. Farahani, *Open J. Math. Anal.*, **2017**, *1*, 44–59. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [30] M.R. Farahani, *Int. J. Theor.l Chem.*, **2013**, *1*, 01-09.

- [31] M.R. Farahani, *Chem. Phys. Res. J.*, **2013**, 6, 27-33.
- [32] M.R. Farahani, W. Gao, *J. Chem. Pharma. Res.*, **2015**, 7, 535-539.
- [33] M.R. Farahani, *J. Chem. Acta*, **2013**, 2, 22-25.
- [34] W. Gao, M.R. Farahani, *J. Nanotech.*, **2016**, 2016, 1-6. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

How to cite this article: Hanan Ahmed, Mohammad Reza Farahani*, Anwar Alwardi, Ruby Salestina M. Domination topological properties of some chemical structures using φ_p -Polynomial approach. *Eurasian Chemical Communications*, 2021, 3(4), 210-218. **Link:** http://www.echemcom.com/article_128247.html