

FULL PAPER

Major flavonoids identification by HPLC-MS/MS and evaluation of antiradical potential of mung bean (*Vigna radiata*) seed extracts in Burkina Faso

Mahamadi Ouedraogo^{a,b,*} | Benjamin Bazie^{a,c} | Dominique Saga Kabore^a | Salfo Ouedraogo^{d,e} | Bazoin Sylvain Raoul Bazie^{f,g} | Elie Kabre^g | Wend-Pagnagde Félicien Marie Serge Zida^h | Benoit Joseph T. Batiéno^h | Moumouni Koala^{a,d} | Eloi Pale^a

^aLaboratoire de Chimie Organique et de Physique Appliquées, Département de Chimie, UFR-SEA, Université Joseph KI-ZERBO, 03 BP 7021 Ouagadougou 03, Burkina Faso

^bUniversité Norbert ZONGO, avce Maurice Yameogo, Koudougou BP 376, Burkina Faso

^cÉcole Polytechnique de Ouagadougou, Ouagadougou, Burkina Faso

^dLaboratoire de Recherche-Développement de Phytomédicaments et Médicaments, Département Médecine, Pharmacopée Traditionnelles et Pharmacie, Institut de Recherche en Science de la Santé (IRSS/CNRST), 03 BP 7047 Ouagadougou, Burkina Faso

^eCentre de Formation, de Recherche et d'Expertise en Sciences du Médicament (CEA-CFOREM), Département sciences et santé, Université Joseph KI-ZERBO, 03 BP 7021 Ouagadougou, Burkina Faso

^fCentre Universitaire de Manga, Département de Biochimie, Manga, Burkina Faso

^gAgence Nationale pour la Sécurité Sanitaire de l'Environnement, de l'Alimentation, du Travail et des produits de santé (ANSSEAT), Département de Biochimie, Ouagadougou, 09 BP Ouagadougou 09, Burkina Faso

^hLaboratoire de Génétique et de Biotechnologies Végétales, Département de Productions Végétales, Institut de l'Environnement et des Recherches Agricoles (INERA) 04 BP 8545 Ouagadougou, Burkina Faso

*Corresponding Author:

Moumouni Koala

E-mail: Moumounikoala@gmail.com

Tel.: +(00226) 76294617

Mung bean is one of the most essential legumes in Burkina Faso due to its short maturation cycle (45 to 55 days) and its nutritional content. This work aims to contribute to a better knowledge of the bioactive compounds of the seeds of Beng Tigré, a mung bean cultivar originating from India and acclimatized in Burkina Faso. To do this, the contents of total phenolic compounds, flavonoids, and carotenoids were determined by colorimetric methods. The mineral elements were assayed by atomic absorption spectrometry after the seeds mineralization. The antiradical potential was evaluated by monitoring the disappearance of the stable radical DPPH• by UV-Visible absorption spectrophotometry at 517 nm and compared to standards. Major flavonoids were identified by high-performance liquid chromatography-tandem mass spectrometry (HPLC-MS/MS). Total phenolic compounds, flavonoids, and carotenoids were 4.356±0.162 mg GAE/g, 3.151±0.262 mg QE/g, and 0.302±0.012 mg BCE/g, respectively. Potassium, magnesium, and calcium were, respectively, the most abundant mineral elements in the seeds: K > Mg > Ca > Fe > Zn > Na > Mn > Cu. An antiradical activity of the Beng Tigré extract was observed with an IC₅₀ of 209.97±13.55 µg of AO/g DPPH•. This activity is less effective than Gallic acid, ascorbic acid, and trolox. Finally, the HPLC-MS/MS analysis identified four compounds: naringenin 8-C-glucoside, luteolin 8-C-glucoside, apigenin 6-C-glucoside, and luteolin 7-O-rutinoside. Due to their high biological potential, these so-called bioactive compounds are the subject of growing research interest.

KEYWORDS

Antiradical potential; bioactive compounds; HPLC-MS/MS; mung bean.

Introduction

The Food and Agriculture Organization of the United Nations (FAO) has declared 2016 the International Year of Pulses [1]. In effect,

legumes are particularly appreciated for their nutrient-rich products, especially the seed and, very often, the vegetable part (leaves and pods) [2]. They are an important source of

proteins, various amino acids, vitamins, and essential mineral elements. These plants also provide so-called bioactive compounds, including, among others, terpenoids, nitrogen compounds, phenolic compounds, etc.

In Burkina Faso, cowpea is the most widespread and consumed legume, alongside which we can mention voandzou, soybeans. However, with climate change, more than the pulses grown in our tropics may be needed to be increased to meet the ever-increasing demand of populations. Thus, the introduction into the Burkina Faso agriculture of mung bean, a legume, could contribute to people's resilience to the effects of climate change by increasing the supply of legumes, achieving food sovereignty, and helping to eradicate malnutrition.

Mung bean is a short-cycle (45-55 days) annual seed legume in the family Fabaceae and is widely grown in tropical and subtropical regions [3]. It is mainly grown in rotation with cereals. Mung bean also represents an opportunity to produce protein-rich foods for rural populations under changing climatic conditions [4-6]. Previous work on mung bean cultivars has shown that the seeds are excellent sources of protein, essential amino acids, minerals, vitamins, and phenolic compounds such as phenolic acids and flavonoids. These chemical substances, with very different physicochemical properties, exhibit various biological activities such as antitumor, antiviral, antifungal, antimicrobial, antioxidant, anti-parasitic, etc. [7]. These compounds are beneficial in preventing and curing many chronic ailments, namely cancer, diabetes, and cardiovascular diseases, due to their antiradical properties [8-10].

Recently, the interest in natural antioxidants has increased considerably in their therapeutic properties [11,12]. Natural antioxidants, the main representatives of which are carotenoids, ascorbic acid, tocopherols, and flavonoids, can protect the human body from free radicals and delay the

progression of many chronic diseases as well as oxidative rancidity of lipids in food [13,14]. Thus, scientific research has been developed in various specialties to extract, identify, and quantify these compounds from several natural medicinal plants and food products.

The present work aims to identify the main flavonoids and evaluate the antiradical potential by assessing the contents of bioactive compounds in the extracts of mung bean seeds (Beng Tigré) popularized in Burkina Faso.

This study provides new scientific information on the mung bean (Beng Tigré) acclimatized and popularized in Burkina Faso. It fills the gap in the antiradical potential and phytochemical profiling of this little-studied variety in the sub-region.

Materials and methods

Plant Materials

The mung bean (*Vigna radiata*) seeds were the subject of our study (Figure 1). They were provided by the plant laboratoire de Génétique et Biotechnologies Végétales de l'Institut de l'Environnement et de Recherche Agricole (INERA) in Kamboinsé (Ouagadougou; Burkina Faso).




FIGURE 1 Mung bean seeds (Beng Tigré)

Chemical products

The chemicals used were: Folin-ciocalteu's reagent (RFC); quercetin; gallic acid; L-ascorbic acid; 2,2-diphenyl-1-picrylhydrazyl (DPPH); ferric 2, 4, 6-tripyridyl-s-triazine (TPTZ); and 6-Hydroxy-2,5,7,8-tetramethyl chroman-2-carboxylic acid (Trolox, a hydrophilic derivative of tocopherol). These products were purchased from Sigma-Aldrich. Ultrapure water (18 M Ω) was prepared with a water purifier system (ELGA PureLab).

All other used chemical reagents were of analytical grade.

Methods

Extraction

The total phenolic compounds of the mung bean were extracted using different types of solvents: the acetone/water/acetic acid system (70:29.5:0.5; v/v/v), ethanol, methanol, and acetone [13,15].

The total carotenoid extracts were obtained from n-hexane, acetone, and an acetone/n-hexane mixture (50:50; v/v) [16]. Now, 1 gram of seed powder was extracted by maceration for 24 hours at 4 °C with 3 mL of each solvent. The extracts were filtered, and the residues were re-extracted twice with 2 mL of solvent for 24 hours. The filtrates are combined and stored at 4 °C to determine the contents of total phenolic compounds, flavonoids, and carotenoids [17].

In addition, the ground seed powder was exhausted by maceration with the acetone/water/acetic acid system (70:29.5:0.5; v/v/v). The filtrate obtained from the extract was concentrated almost to dryness under reduced pressure (T = 40 °C), and then dried in a desiccator under vacuum to have a crude extract which was used for the study of the antiradical activity and identification of flavonoids.

Assay of total phenolic compounds (TPC)

The phenolic compounds were determined by the Folin-Ciocalteu method [18]. This method, initially described by Slinkard and Singleton [19], allows for knowing the total phenolic content of a given sample. Briefly, 60 μ L of the extract was added with 60 μ L of the Folin-Ciocalteu reagent (diluted ten times). After 8 min, 120 μ L of sodium carbonate (Na₂CO₃) was added at 7.5% (w/v). After 30 min of incubation (room temperature), the absorbance was read at 765 nm using a SAFAS-type spectrophotometer. The blank is prepared by replacing the Folin-Ciocalteu reagent with distilled water. A standard curve was established using gallic acid as a standard (Table 1) and the results are expressed in mg of equivalents of gallic acid (GAE) /g of dry seeds.

Assay of total flavonoids (TFT)

Total flavonoid contents were assessed using aluminum trichloride [19]. Briefly, 20 μ L of quercetin with a concentration in the range 0.001 and 0.5 mg/mL or extract was mixed with 10 μ L of NaNO₂ (5%) and left to react for 5 min at room temperature, and then 10 μ L of a solution of AlCl₃ (10%) was added. After 6 min of reaction at ambient temperature, 40 μ L of NaOH (1 M) was added to the reaction medium. The mixture was diluted with 130 μ L of ultrapure water and homogenized. The absorbance was read immediately at 510 nm using a SAFAS-type spectrophotometer.

The total flavonoid content of the extract expressed in mg of quercetin equivalent per gram of dry seeds (mg of QE/g) was obtained by relating the absorbances read on the standard curve established from quercetin (Table 1).

Assay of total carotenoids (TCT)

The total carotenoid contents of the extracts were evaluated according to the method described by McMurry *et al.* [16,20,21] but slightly modified. After suitable dilution, the absorbances of the extracts, kept at room temperature and protected from light, were read at 450 nm.

The total carotenoid contents were obtained by relating the absorbances of the extracts to a standard curve established using β -carotene as a standard (Table 1). The carotenoid contents are expressed in Equivalents of β -carotene per gram of dry seeds (BCE/g).

Assay of mineral elements

The contents of mineral elements were evaluated after the mineralization of the seeds of the Beng Tigré [22]. Indeed, the minerals were subjected to an air-acetylene flame atomic absorption spectrometer of the Analyst PinAAcle 900T type, of the Perkin Elmer brand. The wavelengths of the elements to be analyzed are initially defined on the device (324.75 nm for copper, 213.86 nm for zinc, 248.33 nm for iron, 285.2 nm for magnesium, 766.49 nm for potassium, 279.48 nm for manganese, and 589.0 nm for sodium). Next, the different readings of the calibration ranges made it possible to establish the calibration curves reflecting the absorbance as a function of the concentration. Finally, the samples to be analyzed are presented to the apparatus to determine their absorbances [23]. A blank is obligatorily passed between the passages of two different solutions. After the analysis of ten (10) samples, a control solution is passed to verify the effectiveness of the analysis results.

Anti-radical activity by the DPPH• radical test

The free radical scavenging activity of the seed extracts was measured using the method described by Burits and Bucar [24], with some modifications. The commercial radical of DPPH• was dissolved in methanol at a concentration of 0.2 mg/mL and kept at 4 °C protected from light. 50 μ L of the crude extract solution dissolved in methanol at different concentrations ranging from 0.875 mg/mL to 6.8×10^{-3} mg/mL were mixed with 200 μ L of a DPPH• solution. The absorbance was measured at the wavelength of 517 nm every 30 seconds for 30 min. The inhibition of the DPPH• free radical by each standard used (ascorbic acid (vitamin C), gallic acid, and Trolox) was also analyzed at the same concentration to make the comparison. Antioxidants, with the property of donating hydrogen to the DPPH• radical lead to the discoloration of DPPH solution, which was taken on a color ranging from dark pink to pale yellow depending on the antioxidant content (Figure 2). The percentage of the DPPH• radical remaining after the radical reaction carried out by an antioxidant was calculated according to the formula [25]:

$$\%DPPH \bullet_{Rest} = \frac{Abs_{DPPH+antioxydant}}{Abs_{DPPH(T=0)}} \times 100$$

$\%DPPH \bullet_{Rest}$ is proportional to the concentrations of the antioxidant. The concentration that causes a decrease of the initial concentration of DPPH• by 50% was IC_{50} . This parameter is inversely proportional to the antioxidant capacity and is obtained graphically from the curve relating the percentage reduction $\%DPPH \bullet_{Rest}$ and the concentration of antioxidants. It can be further expressed in anti-radical effectiveness (EAR = $1/IC_{50}$).

the acetone/n-hexane system extract with 0.302 ± 0.012 mg of BCE/g of dry seeds, followed by the acetone extract (0.183 ± 0.014 mg BCE/g) and n-hexane extract (0.054 ± 0.002 mg BCE/g). The carotenoid extracts can be classified according to this order: acetone/n-hexane > acetone > n-hexane. The results show that the acetone/water/acetic acid system is the most appropriate solvent for extracting total phenolic compounds in mung bean seeds. This confirms the works of the literature, where this solvent system was unanimously used for the total phenolic compounds extraction [16]. Moreover, with this solvent system, the contents of total phenolic compounds (4.356 ± 0.162 mg of GAE/g) and flavonoids (3.151 ± 0.262 mg of QE/g) are comparable to those of Voandzou (≤ 4.536 mg GAE/g; 4.282 ± 0.093 mg QE/g) and cowpea (5.584 ± 0.037 mg GAE/g; 4.558 ± 0.156 mg QE/g) [26]. The total carotenoid assay results

showed that the acetone/n-hexane system extracts carotenoids better. This could be explained by the fact that each solvent extracts a specific group of carotenoids. In fact, polar or slightly polar molecules, particularly xanthophylls, are soluble in acetone, and the group of apolar carotenes is more soluble in hexane. Thus, the acetone/n-hexane system extracts both xanthophylls and carotenes.

These results agree with several authors who revealed that mixed solvents efficiently extract bioactive compounds such as phenolic compounds and carotenoids [28-30]. Indeed, using mixed solvents results in a strong enrichment of the extracts in bioactive compounds. The advantage of composite solvents would be due to the increase in the solubility of bioactive compounds in the extracts obtained by these solvents compared to those obtained by pure solvents [29].

TABLE 1 Establishment of standard curves

Standard curves	Standards	Calibration equation	Correlation coefficient
CPT	Gallic acid	$y = 17,814x + 0,0485$	$R^2 = 0,9997$
TFT	Quercetin	$y = 0,767x + 0,0862$	$R^2 = 0,9963$
TCT	β -carotene	$y = 11,618x - 0,0075$	$R^2 = 0,9994$

Mineral elements content

The mineral element contents of the Beng tigré seeds are presented in Table 2. The results make it possible to distinguish the elements present in significant quantities, including Potassium (1140.2 ± 14.851 mg/100 g),

Magnesium (146.326 ± 1.912 mg/ 100 g), Calcium (51.529 ± 0.644 mg/100g), and those in minute quantities such as Iron (21.96 ± 0.707 mg/100 g), Zinc (18.697 ± 0.568 mg/100 g), Manganese (4.22 ± 0.098 mg/100 g) and Copper (1.275 ± 0.052 mg/100 g).

TABLE 2 Mineral element content of Beng Tigré seeds

Mineral elements	Contents (mg/100g)	PRI (mg/day)	SAM
Cu	1.275±0.052	1.1	0.6
Zn	18.697±0.568	7-9.5	4-5
Mn	4.22±0.098	1-10	-
Fe	21.96±0.707	1.5	-
Na	12.32±0.041	600-3500	-
K	1140.2±14.851	3100	1600
Ca	51.529±0.644	700	400
Mg	146.326±1.912	150-500	-

PRI: Population Reference Intake and SAM: Minimum Intake Threshold.

Sodium and potassium play the role of regulating the water content of human body and participate in maintaining the acid-base balance. The Na/K ratio in the seeds of the Beng tigré is equal to 0.01, less than 1. It is deduced that consuming this legume is beneficial in the fight against cardiovascular diseases [31,32]. Zinc and manganese are antioxidant trace elements that promote the elimination of excess free radicals in the body. Moreover, 100 g of Beng tabby learned appears sufficient to cover the daily requirements of iron, zinc, manganese, and copper. In any case, 200 g of Beng tigré covers the minimum intake threshold.

The mineral elements in our bodies play a crucial role in our daily lives. They compensate for daily losses (perspiration, urine, stools) and satisfy needs in certain physiological situations (bone growth, pregnancy, breastfeeding, and menstruation).

Study of antiradical activity

Antioxidant activity defines the ability of a compound to protect the body against the aggressiveness of free radicals. Its determination by the DPPH test is one of the methods for biologically and nutritionally evaluating food quality [33]. In addition, this test makes it possible to assess the kinetics of



trapping free radicals. Indeed, in the presence of a hydrogen donor antioxidant, the free radical DPPH• is reduced to a stable molecule DPPH₂. This causes a decrease in the DPPH• concentration and a reduction in absorbance during the reaction time until the hydrogen-donating antioxidant capacity is exhausted. Thus, at different concentrations of the antioxidant tested, the kinetics of reduction of mDPPH• is monitored over time until an equilibrium is obtained, reflected by the presence of a plateau at the equilibrium time (Teq). The percentage of DPPH• remaining (%DPPH•_{Rest}) also decreases until it reaches a plateau. This kinetic varies according to the antioxidant used. The results are depicted in Figure 2. The reaction between DPPH and the reference hydrogen-donating antioxidants (gallic acid, ascorbic acid, and trolox) reaches equilibrium after a short time compared to the mung bean seed extract (Table 3). Indeed, the time required to reach equilibrium varies according to the concentrations. For gallic acid, equilibrium is reached in 30 seconds for a concentration of 54.69 µg/mL and 1 min 30 s for 6.84 µg/mL and 3.42 µg/mL. The equilibrium time of ascorbic acid is 30 seconds for the different concentrations tested (54.69 µg/mL, 6.84 µg/mL, and 3.42 µg/mL). Trolox has an equilibrium time of 1 minutes and 30

seconds in 13.67 $\mu\text{g}/\text{mL}$ and 2 minutes in 6.84 $\mu\text{g}/\text{mL}$ and 3.42 $\mu\text{g}/\text{mL}$. The equilibrium time observed for the mung bean seed extract is 1 minute and 30 seconds. The amount of antioxidants required to decrease the concentration of the free radical DPPH \bullet by 50% (IC_{50}) was calculated from the equation of the curve of the percentage reduction of DPPH \bullet as a function of the antioxidant concentration (Figure 3). The anti-radical efficacy EAR is deduced by the inverse of the IC_{50} ($\text{EAR}=1/\text{IC}_{50}$). The IC_{50} values of gallic acid, ascorbic acid, trolox, and mung bean seed extract are presented in Table 3. Statistical analysis of IC_{50} and EAR showed no significant difference between gallic acid, ascorbic acid, and Trolox. These antioxidants have similar efficacy but are more active than the mung bean seed extract, which restores the stability of DPPH with an IC_{50} of 209.97 ± 13.55 $\mu\text{g}/\text{g}$.

According to these results, it is proven that the reference antioxidants used (ascorbic acid, gallic acid, and trolox) are more effective compared to the extract of mung bean seeds. However, it should be noted that mung bean seed extract is a crude extract that contains bioactive compounds such as phenolics, flavonoids, carotenoids, mineral elements, etc.

Identification of mung bean flavonoids by HPLC-MS/MS

The crude extract of the Beng Tigré subjected to HPLC-MS made it possible to observe several significant signals (Figure 4) corresponding to protonated molecules $[\text{M}+\text{H}]^+$. Four of these signals [**A**: $t_{\text{R}}=37.64$ min, **D**: $t_{\text{R}}=47.32$ min, **E**: $t_{\text{R}}=58.35$ min, and **F**: $t_{\text{R}}=64.67$ min] likely to correspond to flavonoids were subjected to fragmentation in MS/MS.






FIGURE 3 Kinetics of reduction of DPPH• by antioxidants

TABLE 3 Characteristic parameters of the DPPH• radical reduction kinetics

	Concentration (µg/mL)	Teq (min)	%DPPH _R at equilibrium	Curve equation	Correlation coefficient	IC ₅₀ (µg/g)	IC ₅₀ (mean±standard deviation)	EAR (mean±standard deviation)
Gallic acid	54.69	0.5	15.84158416	$y = 3.3368x^{-0.532}$	$R^2 = 0.996$	6.17		
	6.84	1.5	39.24422442	$y = 3.4377x^{-0.518}$	$R^2 = 0.989$	5.69	6.17±0.43 ^a	1.75.10 ⁻¹ ±0.013 ^a
	3.42	1.5	68.31683168	$y = 3.2485x^{-0.522}$	$R^2 = 0.983$	5.32		
Ascorbic acid	54.69	0.5	3.630363036	$y = 0.1721x^{-1.064}$	$R^2 = 0.990$	4.84		
	6.84	0.5	40.92409241	$y = 0.1729x^{-1.062}$	$R^2 = 0.990$	4.79	4.80±0.04 ^a	2.08.10 ⁻¹ ±0.002 ^a
	3.42	0.5	63.69636964	$y = 0.1435x^{-1.095}$	$R^2 = 0.989$	4.77		
Trolox	13.67	1.5	21.78217822	$y = 5.7838x^{-0.445}$	$R^2 = 0.987$	7.8		
	6.84	2	48.18481848	$y = 3.4965x^{-0.531}$	$R^2 = 0.985$	6.6	6.79±0.93 ^a	1.49.10 ⁻¹ ±0.02 ^a
	3.42	2	67.98679868	$y = 2.0351x^{-0.625}$	$R^2 = 0.979$	5.98		
Extract (mung bean seeds)	880	1.5	6.270627063	$y = 6.6904x^{-1.342}$	$R^2 = 0.933$	223.4		
	440	5.5	9.900990099	$y = 6.2503x^{-1.333}$	$R^2 = 0.949$	210.2	209.97±13.55 ^b	31.10 ⁻⁵ ±0.005 ^b

IC₅₀ and ERA results are expressed as mean ± standard deviation. Values are processed by IBM SPSS Statistics 25.0 software. The significant difference between the values is indicated by letters ($P < 0.05$).




FIGURE 4 Percentage reduction of the DPPH• radical according to the concentration of the antioxidant

From these signals, the following compounds were detected: compound 1 at m/z 435 (**D** signal), compound 2 at m/z 449 (**A** signal), compound 3 at m/z 595 (**E** signal), and compound 4 at m/z 433 (**F** signal). These compounds corresponded to the raw formulas' calculated masses $\text{C}_{21}\text{H}_{23}\text{O}_{10}$, $\text{C}_{21}\text{H}_{21}\text{O}_{11}$, $\text{C}_{27}\text{H}_{31}\text{O}_{15}$, and $\text{C}_{21}\text{H}_{21}\text{O}_{10}$, respectively. These formulas can also check to those of flavonoids of the monoglycosylated (mono-*O*-glycosylated or mono-*C*-glycosylated) or diglycosylated flavone type. A previous study found that the most abundant flavonoids in mung bean possess the aglycone of apigenin or luteolin [34]. Furthermore, it should be remembered that the neutral loss of 132, 162, and 146 atomic mass units (u) make it possible to identify pentosides (xylose or arabinose), hexosides

(glucose or galactose), and deoxyhexoside (rhamnose), respectively, in the case of *O*-glycosylated flavonoids [35]. Regarding *C*-glycosylated flavonoids, in negative ESI, the observation of a neutral loss of 120 u and 90 u corresponds to a hexose pattern, a loss of 104 u and 74 u corresponds to a deoxyhexose pattern, and a loss of 90 u and 60 u to that of a pentose motif. In positive ESI, the loss of 120 u shows the presence of a hexose [36]. Thus, analysis of the MS/MS spectrum of compound 1 m/z 435 (Figure 5) shows the presence of fragment ions characteristic of *C*-glycosylated flavonoids. Indeed, the presence of the ion at m/z 315 ($[\text{O}^2\text{X}]^+$) commonly encountered among the fragment ions of *C*-glycosylated flavonoids and dependent on the presence of a hexose (glucose or galactose) is observed. This fragment ion, by loss of 42 u, makes it

possible to deduce the nature of the aglycone of the flavonoid which is here naringenin ($[^{0,2}X-42]^+$ [25]). According to the literature, the *C*-glycosylation of flavonoids is carried out in position 6 or 8 of the aglycone [37]. Therefore, the presence or absence of certain fragment ions in the MS/MS spectrum makes it possible to distinguish the 8-*C* isomers from the 6-*C* glycosides [38]. In the case of 6-*C* glycosides, the ion at m/z $[M+H-120 ; ^{0,2}X]^+$ fragments in turn to give, for example, the m/z ions $[^{0,2}X-CHO]^+$ and $[^{0,2}X-H_2O]^+$. These ions could not be observed in the fragmentation of compound 1. In addition, there is often a fragmentation of the *C* cycle in the 6-*C* glycosides, thus giving fragment ions at m/z $[^{1,3}A]^+$ and $[^{0,2}B]^+$. These ions are also not observed. Therefore, compound 1 would likely be naringenin 8-*C*-glycoside. Since glucose is more common than galactose, compound 1 would be naringenin 8-*C*-glucoside. Compound 2 detected at m/z 449 corresponds to the molecule eluted after 37.64 min. The corresponding CID (Collision Induced Dissociation) spectrum is presented in Figure 6. On CID, the molecule undergoes mainly neutral losses of 120 u and 150 u leading to fragment ions at m/z 329 $[M+H-C_4H_8O_4]^+$ and 299 $[M+H-C_4H_8O_4-CH_2O]^+$, respectively. As mentioned above, these losses are compatible with the characteristic fragment ions of a *C*-glycoside indicating the presence of a hexose (glucose) linked to a carbon of the genin. The 6-*C* and 8-*C* positions of the sugar attached to the genin in the *C*-glycosylated flavonoids can be determined by observing the intensity of the ion corresponding to the first loss of water [38]. In ESI+, if the ion $[M+H-H_2O]^+$ is the most abundant fragment peak of the MS/MS spectrum, the sugar is linked to carbon 8 of the genin. Otherwise, it is in position 6-*C* [25]. Also, the peak of 329 m/z is more abundant than that at 299 m/z in the case of an 8-*C*. The opposite is observed in the case of a 6-*C* [36]. These observations thus make it possible to deduce that compound 2 is probably luteolin

8-*C*-glucoside. Compound 3 (m/z 595) eluted after 58.35 min undergoes two consecutive losses of 146 u and 162 u, revealing the presence of a hexose residue (glucose) and a deoxyhexose residue (rhamnose) attached to the aglycone group. On the CID spectrum (Figure 7), the presence of the peak at 449 m/z ($[M+H-146]^+$) implies that one sugar unit is bound to another (osidic bond) and not directly to the aglycone [39]. This indicates an *O*-diglycosylated structure. A peak at 287 m/z ($[M+H-146-162]^+$) corresponding to the loss of a rutinoside is also observed. This peak at 287 m/z would correspond to the aglycone of luteolin or Kaempferol. A study showed that luteolin is more common in mung bean seeds [34]. Literature data also show that flavonoids could be glycosylated in any position, but *O*-glycosylation mainly occurs in position 7, as in flavones, isoflavones, flavanones, and flavonols [40,41]. Given these analyses, Compound 3 squeezed luteolin 7-*O*-rutinoside. Compound 4 eluted after 64.67 min corresponds to a compound detected at m/z 433. The MS/MS spectrum analysis does not show any characteristic fragment of the sugar loss (Figure 8). The sugar would therefore be linked to the aglycone by a carbon-carbon bond (*C*-glycosylation). Fragment ion analysis shows ions at m/z 283 $[^{0,2}X-CH_2O]^+$ and 313 $[^{0,2}X]^+$, which are characteristic of *C*-glycosylated flavonoids [42]. Peaks at m/z 270 $[^{0,1}X]^+$, 283 $[^{0,2}X-CH_2O]^+$ would be specific to 6-*C*-glycoside [37]. These fragment ions are observed on the MS/MS spectrum of compound 4. In addition, the fragment ions 415, 397, and 379 m/z corresponding to respective losses of one, two, and three molecules of H_2O are characteristic fragmentation pathways vitexin (Apigenin 8-*C*-glycosylated) or isovitexin (Apigenin 6-*C*-glycosylated) [43]. Considering all these results and the data from the literature, the probable structure of compound 4 is 6-*C*-glycosylated apigenin. Today, many techniques are employed for the structural analysis of bioactive compounds.

These include UV-Visible spectroscopy, nuclear magnetic resonance spectroscopy, infrared spectroscopy, Raman spectroscopy, mass spectrometry, and X-ray crystallography. However, there is yet to be a universal analytical technique for analyzing chemical substances. Each method has its advantages and disadvantages. The application of one of these techniques will depend on several parameters, including the nature of the sample to be analyzed and the

study's objective of the problem being addressed. These techniques are complementary and can be combined in different ways. This is the case of the HPLC-MS/MS combination used in this study. This technique is advantageous because it identifies non-volatile raw extract molecules without isolation steps. Moreover, although expensive, it is a reference technique for its flexibility, sensitivity, specificity, and speed.








FIGURE 5 HPLC chromatogram of crude mung bean extract




Chromatogram of D signal


**FIGURE 6** ESI+-MS/MS spectrum of compound **1** (m/z 435)

Chromatogram of A signal

**FIGURE 7** ESI+-MS/MS spectrum of compound **2** (m/z 449)



Chromatogram of E signal

FIGURE 8 ESI+-MS/MS spectrum of compound **3** (m/z 595)

Conclusion

In this work, we analyzed the bioactive compounds and their antiradical activity, and also identified some flavonoids of the seeds of Beng Tigré to enhance this legume and promote its consumption. This study showed that Beng Tigré is a potential source of phenolic compounds, flavonoids, carotenoids, and mineral salts. According to the results, the crude extract using acetone/water/acetic acid solvent system of the seeds showed significant antiradical activity ($IC_{50} =$

$209.97 \pm 13.55 \mu\text{g/g}$). This indicates that the culture of Beng Tigré is vital in nutrition, and its regular consumption naturally contributes to preventing or curing specific chronic ailments such as cardiovascular diseases, cancers, diabetes, etc. The major molecules of flavonoids identified, namely naringenin 8-C-glucoside, luteolin 8-C-glucoside, apigenin 6-C-glucoside, and luteolin 7-O-rutinoside also show that Beng Tigré could be a source natural supply of colorants or food additives, pharmaceutical and cosmetic active

ingredients, to the pharmaceutical, cosmetic and food industries.

Acknowledgements

This research study was made possible thanks to the technical support of the laboratory de Génétique et Biotechnologies Végétales de l'Institut de l'Environnement et des Recherches Agricoles (INERA) of Kamboinsé (Ouagadougou; Burkina Faso) and Agence Nationale pour la Sécurité Sanitaire de l'Environnement, de l'Alimentation, du Travail et des produits de santé (ANSSEAT-Ex Laboratoire National de Santé Publique) of Burkina Faso. The authors thus thank the staff of INERA for the supply of plant material and of the ANSSEAT for its contribution to the acquisition of HPLC-MS/MS.

Funding

This work was funded by Centre de

Formation, de Recherche et d'Expertise en Sciences du Médicament (CEA-CFOREM), l'école doctorale sciences et santé, Université Joseph KI-ZERBO.

Conflict of Interest

There are no competing interests to declare that are relevant to the content of this article.

Orcid:

Mahamadi Ouedraogo:

<https://orcid.org/0009-0002-2733-9031>

Bazoin Sylvain Raoul Bazie:

<https://orcid.org/0000-0001-8225-2217>

Elie Kabre:

<https://orcid.org/0000-0003-4406-6354>

Wend-Pagnagde Félicien Marie Serge Zida:

<https://orcid.org/0009-0006-9389-4091>

Moumouni Koala:

<https://orcid.org/0000-0001-7241-9183>

References

- [1] S. Snapp, M. Rahmanian, C. Batello, T. Calls, Légumes secs et exploitations durables en Afrique subsaharienne, *FAO.*, **2018**, 68. [[Google Scholar](#)], [[Publisher](#)]
- [2] R.A. Dixon, L.W. Sumner, Legume natural products: understanding and manipulating complex pathways for human and animal health, *Plant Physiol.*, **2013**, *131*, 878-885. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [3] X. Peng, Z. Zheng, K.W. Cheng, F. Shan, G.X. Ren, F. Chen, M. Wang, Inhibitory effect of mung bean extract and its constituents vitexin and isovitexin on the formation of advanced glycation endproducts, *Food Chem.*, **2008**, *106*, 475-481. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [4] J.D.H. Keatinge, W.J. Easdown, R.Y. Yang, M.L. Chadha, S. Shanmugasundaram, Overcoming chronic malnutrition in a future warming world: the key importance of mungbean and vegetable soybean, *Euphytica.*, **2011**, *180*, 129-141. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [5] J.E. Arsenault, R.J. Hijmans, K.H. Brown, Improving nutrition security through agriculture: an analytical framework based on national food balance sheets to estimate nutritional adequacy of food supplies, *Food Secur.*, **2015**, *7*, 693-707. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [6] C.H. Foyer, H.M. Lam, H.T. Nguyen, K.H. Siddique, R.K. Varshney, T.D. Colmer, W. Cowling, H. Bramley, T.A. Mori, J.M. Hodgson, J.W. Cooper, Neglecting legumes has compromised human health and sustainable food production, *Nat. Plants*, **2016**, *2*, 16112. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [7] M. Khalil, S. Noor, Z. Ahmad, F. Ahmad, Fate of Pakistani exported Mango due to its toxicity (heavy metals, pesticides, and other toxic organic components), *J. Appl. Organomet. Chem.*, **2023**, *3*, 86-107. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

- [8] Y. Yao, X. Yang, J. Tian, C. Liu, X. Cheng, G. Ren, Antioxidant and antidiabetic activities of black mung bean (*Vigna radiata* L.), **2013**, *J. Agric. Food Chem.*, **61**, 8104–8109. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [9] J. Luo, W. Cai, T.B. Wua, B. Xu, Phytochemical distribution in hull and cotyledon of adzuki bean (*Vigna angularis* L.) and mung bean (*Vigna radiata* L.), and their contribution to antioxidant, anti-inflammatory and anti-diabetic activities, *Food Chem.*, **2016**, **201**, 350-360. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [10] J.P. Singh, A. Kaur, N. Singh, L. Nim, K. Shevkani, H. Kaur, D.S. Arora, In vitro antioxidant and antimicrobial properties of jambolan (*Syzygium cumini*) fruit polyphenols, *LWT-Food Sci. Technol.*, **2016**, **65**, 1025-1030. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [11] H.H. Hama, I. Moussa, K. Ikhiri, B. Ouedraogo, R. Adama, Activité antioxydante des extraits méthanoliques de différents organes de *Detarium microcarpum*, *European Sci. J.*, **2019**, **15**, 1857-7881. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [12] G. Lal Gupta, N. Patil Samant, Acute and sub-acute oral toxicity evaluation of avicularin, *J. Med. Chem. Sci.*, **2023**, **6**, 2327-2337. [[crossref](#)], [[Pdf](#)], [[Publisher](#)]
- [13] L.S. Lai, S.T. Chou, W.W. Chao, Etudes sur l'antioxydantes idatives de *Hsian-tsoa* (*Mesona procumbens* Hemsl) feuille de gomme, *J. Agri. Food Chem.*, **2001**, **49**, 963-968. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [14] N.A. Patil, S. Udgire, D.R. Shinde, P.D. Patil, Green synthesis of gold nanoparticles using extract of *Vitis vinifera*, *Buchananialanza*, *Juglandaceae*, *Phoenix Dactylifera* plants, and evaluation of antimicrobial activity, *Chem. Methodol.*, **2023**, **7**, 15-27. [[crossref](#)], [[Pdf](#)], [[Publisher](#)]
- [15] R.E. Wrolstad, T.E. Acree, E.A. Decker, M.H. Penner, D.S. Reid, S.J. Schwartz, C.F. Shoemaker, D. Smith, P. Sporns, Handbook of food analytical chemistry pigments, colorants, flavors, texture, and bioactive food components, *John Wiley and Sons, Inc.*, **2005**, 81-90. [[Google Scholar](#)], [[Publisher](#)]
- [16] A.H. Moumouni Koala, K. Somé, E. Palé, A. Sérémé, J. Belem, M. Nacro, Evaluation of Eight Orange Fleshed Sweetpotato (OFSP) Varieties for Their Total Antioxidant, Total Carotenoid and Polyphenolic Contents, *J. Nat. Sci. Res.*, **2013**, **3**, 67-72. [[Google Scholar](#)], [[Publisher](#)]
- [17] O. Mahamadi, H. Adama, W.F.M. Serge, J.T. Benoit, P. Eloi, N. Mouhoussine, Détermination des paramètres physico-chimiques et des teneurs en micronutriments antioxydants de dix variétés de mung bean (*Vigna radiata*) produits dans les conditions agroécologiques du Burkina Faso, *J. Chem. Biol. Phy. Sci.*, **2021**, **11**, 254-274. [[crossref](#)], [[Pdf](#)], [[Publisher](#)]
- [18] G. Chala, Review on green synthesis of iron-based nanoparticles for environmental applications, *J. Chem. Rev.*, **2023**, **5**, 1-14. [[crossref](#)], [[Pdf](#)], [[Publisher](#)]
- [19] A.R. Maleki, L. Nateghi, P. Rajai, The Effect of microwave pretreatment on the extraction rate of flavonoid, anthocyanins, antioxidant compounds and antimicrobial activity of *Punica Granatum* Var. *Pleniflora* (Persian Golnar), *Chem. Methodol.*, **2022**, **6**, 280-292. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [20] Y. Jun, F. Lingling, X. Jian, X. Yedan, Ultrasound-assisted extraction of corn carotenoids in ethanol, *Int. J. Food Sci. Technol.*, **2011**, **46**, 2131-2131. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [21] J. Mc Murry, Organic Chemistry, 7th Edition, Thomson Brooks/ Cole., **2008**, 504 chapter 14.
- [22] P. Sáez-Plaza, M.J. Navas, S. Wybraniec, T. Michałowski, A.G. Asuero, An overview of the Kjeldahl method of nitrogen determination. part II. sample preparation, working scale, instrumental finish, and quality control, *Crit. Rev. Anal. Chem.*, **2013**, **43**, 224-272. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [23] B. Ou, R.L. Prior, D. Huang, The chemistry behind dietary antioxidant capacity assays, *J. Agri. Food Chem.*, **2005**, **53**, 1841-1856. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

- [24] A. Hatami, Preparation, Description and evaluation of the lethality acid loaded liposomal nanoparticles against in vitro colon and liver cancer, *J. Chem. Rev.*, **2021**, *3*, 121-133. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [25] M. Ouedraogo, A. Hema, B.S.R. Bazié, S.W. Zida, B.J.T. Batiéno, E. Kabré, E. Palé, M. Nacro, Identification of major flavonoids using HPLC-MS/MS and determination of antioxidant potential of *Vigna radiata* seed extracts, *Journal de la Société Ouest-Africaine de Chimie.*, **2021**, *050* 57-67. [[Google Scholar](#)], [[Publisher](#)]
- [26] A. Mbaïogaou, A. Hema, M. Ouedraogo, E. Palé, M. Naitormbaide, Y. Mahamout, M. Nacro, Etude comparative des teneurs en polyphénols et en antioxydants totaux d'extraits de graines de 44 variétés de voandzou (*Vigna subterranea* (L.)Verdcourt), *Int. J. Biol. Chem. Sci.*, **2013**, *7*, 861-871. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [27] N. Parisa, M.T. Kamaluddin, M.I. Saleh, E.S. Sinaga, R. Umi Partan, I. Irfannuddin, Flavonoids as Antioxidants: A Review on Tempuyung Plant (*Sonchus Arvensis*), *J. Med. Chem. Sci.*, **2023**, *6*, 2310-2318. [[crossref](#)], [[Pdf](#)], [[Publisher](#)]
- [28] Z. Mohammedi, F. Atik, Impact of solvent extraction type on total polyphenols content and biological activity from *Tamarix aphylla* (L.) karst, *Int. J. Pharma. Bio. Sci.*, **2011**, *2*, 609-615. [[Google Scholar](#)], [[Publisher](#)]
- [29] N. Trabelsi, W. Megdiche, R. Ksouri, H. Falleh, S. Oueslati, B. Soumaya, H. Hajlaoui, C. Abdelly, Solvent effects on phenolic contents and biological activities of the halophyte *Limoniastrum monopetalum* leaves, *LWT-Food Sci. Technol.*, **2010**, *43*, 632-639. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [30] F.I. Ahmadi, R. Fathollahi, D. Dastan, Phytochemical constituents and biological properties of *Scutellaria Condensata* Subsp, *Pycnotricha*, *J. Appl. Organomet. Chem.*, **2022**, *2*, 119-128. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [31] F.J. He, G.A. Macgregor, Beneficial effects of potassium on human health, *Physiol Plant.*, **2008**, *133*, 725-735. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [32] V.E. Kemi, M. Kärkkäinen, C.J. Lamberg-Allardt, High phosphorus intakes acutely and negatively affect Ca and bone metabolism in a dose-dependent manner in healthy young females, *British J. Nut.*, **2006**, *96*, 545-552. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [33] J. Sochor, M. Ryvolova, O. Krystofova, P. Salas, J. Hubalek, V. Adam, L. Trnkova, L. Havel, M. Beklova, J. Zehnalek, I. Provaznik, Fully automated spectrometric protocols for determination of antioxidant activity: advantages and disadvantages, *Molécules*, **2010**, *15*, 8618-8640. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [34] D. Cao, H. Li, J. Yi, J. Zhang, H. Che, J. Cao, L. Yang, C. Zhu, W. Jiang, Antioxidant properties of the mung bean flavonoids on alleviating heat stress, *PLoS. One*, **2011**, *6*, 2010-2071. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [35] L.L. Saldanha, W. Vilegas, A.L. Dokkedal, Caractérisation des flavonoïdes et des acides phénoliques dans *Myrcia Bella* cambess. Utilisation de FIA-ESI-IT-MSn et HPLC-PAD-ESI-IT-MS combinés à la RMN, *Molécules.*, **2013**, *18*, 8402-8416. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [36] M.F. Lubis, V.E. Kaban, K. Gurning, P. Parhan, H. Syahputra, N.A. Juwita, R. Astyka, I. Zulfansyah, Phytochemicals and biological activities of ethanolic extract of *Garcinia atroviridis* leaf grown in indonesia, *J. Med. Chem. Sci.*, **2023**, *6*, 2456-2469. [[crossref](#)], [[Pdf](#)], [[Publisher](#)]
- [37] F. Laëtitia, Signature moléculaire de milieux complexes : Stratégie de couplage à la spectrométrie de masse et interprétation des données, Thèse de Doctorat. Université d'Orléans, **2019**, 504-674. [[Google Scholar](#)], [[Publisher](#)]
- [38] W. Patrice, W. Jean-Luc, N. Karine, R.H. Kirsten, J.M. Hilary, H. Kurt, Evaluation of quadrupole time-of-flight tandem mass spectrometry and ion-trap multiple-stage mass spectrometry for the differentiation of

Cglycosidic flavonoid isomers, *J. Chromatogr. A.*, **2001**, 926, 29-41. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

[39] V. Vukics, V. Guttman, Caractérisation structurale des glycosides flavonoïdes par spectromètre de masse à plusieurs étages y, *Mass Spectrometry Reviews*, **2010**, 29, 1-16. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

[40] F. Cuyckens, Y.L. Ma, G. Pocsfalvi, M. Claeys, Stratégies spectrales de masse en tandem pour la caractérisation structurale des glycosides flavonoïdes, *Analusis.*, **2000**, 28, 888-895. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

[41] W.O. Jean Claude, K. Moumouni, O. Noufou, B.K. Félix, G. Pascal, L.B.C. Yvonne, Total phenolics and total flavonoid contents, antioxidant activity and flavonoids identification by high-performance liquid chromatography-tandem mass spectrometry of *odontonema strictum* (Acanthaceae) leaves, *Asian J. Plant Sci. Res.*, **2017**, 7, 54-63.

[42] E. Chan, S. Yap, A. Lau, P. Leow, H. Koh, Ultra-performance liquid chromatography/time-of-flight mass spectrometry based

metabolomics of raw and steamed Panax notoginseng, *Rapid Commun. Mass Spectrom.*, **2007**, 21, 519-528. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

[43] B. Abad-García, S. Garmón-Lobato, L.A. Berrueta, B. Gallo, F. Vicente, New features on the fragmentation and differentiation of C-glycosidic flavone isomers by positive electrospray ionization and triple quadrupole mass spectrometry, *Rapid Commun. Mass Spectrom.*, **2008**, 22, 1834-1842. [[crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

How to cite this article: Mahamadi Ouedraogo, Benjamin Bazie, Dominique Saga Kabore, Salfo Ouedraogo, Bazoin Sylvain Raoul Bazie, Elie Kabre, Wend-Pagnagde Félicien Marie Serge Zida, Benoit Joseph T. Batiéno, Moumouni Koala, Eloi Pale. Major flavonoids identification by HPLC-MS/MS and evaluation of antiradical potential of mung bean (*Vigna radiata*) seed extracts in Burkina Faso. *Journal of Medicinal and Pharmaceutical Chemistry Research*, 2023, 5(10), 935-952. **Link:** http://jmpcr.samipubco.com/article_177587.html