

**FULL PAPER**

# Differences in anthropometric status in late preterm babies fed with breastmilk, infant milk, and a combination of both

Indana Lazulfa  | Martono Tri Utomo\*  | Nur Aisiyah Widjaja 

Department of Child Health, Faculty of Medicine  
Universitas Airlangga, Dr. Soetomo General  
Academic Hospital, Surabaya, East Java,  
Indonesia

Breast milk is recommended to provide to every baby, including late preterm babies. However, some parents or caregivers choose alternative nutrition over breast milk for various reasons. This study aimed to analyze the differences in body weight, body length, and head circumference of late preterm babies at the ages of 1, 3, and 6 months fed with breast milk, infant milk, and a combination of both. It is a retrospective study. The subjects were children aged 6-12 months born at a gestational age of 34 - <37 weeks, who came to the pediatric outpatient and inpatient clinic at Dr. Soetomo General Hospital between March and June 2023. We used a questionnaire to interview the parents or primary caregivers. One-way ANOVA test was used for normally distributed data and Kruskal-Wallis for not normally distributed data. The sample of this study consisted of 70 respondents out of 106. There was no significant difference in the gender. Most subjects were born at 36 -<37 weeks of gestation (41%). The majority of subjects were raised by mothers (94.3%). The statistical analysis showed that late preterm babies who were fed breast milk, infant milk, and a combination of both did not differ significantly in body weight, body length, and head circumference at the ages of 1, 3, and 6 months. No difference was found in the anthropometric status of late preterm babies at Dr. Soetomo General Hospital who were fed with breast milk, infant milk formula, and a combination of both.

**\*Corresponding Author:**

Martono Tri Utomo

Email: [martono.utomo73@gmail.com](mailto:martono.utomo73@gmail.com)

Tel.: +62 817-0366-7063

**KEYWORDS**

Late preterm; infant nutrition; anthropometric status; breast milk.

**Introduction**

Late preterm babies are unique because, while some physically resemble full-term babies, they face four times the risk of medical complications [1,2]. Some morbidity that can

be found in late preterm babies include asphyxia, respiratory distress, hypothermia, hypoglycemia, hyperbilirubinemia, and sepsis [3]. This can be caused by the immaturity of certain body organs or body functions. It can also be related to immaturity of oromotor

development [4-7]. Early in life, they often require medical care and may receive parenteral nutrition or infant formula milk. However, these babies need breast milk to support their immunity [8-10]. In Indonesia, breastfeeding rate was increased from 30.2% in 2013 to 74.5% [11]. After being discharged from the hospital, some babies still do not receive breast milk for any reason based on maternal, baby, and environmental factors (e.g., accessibility, nipple confusion, low milk production, etc) [12-14]. Infant formula milk can be given as the main nutrition for babies if breast milk is inadequate. It has a standardized composition that approaches the nutrition of breast milk as stated in The Codex Alimentarius, which was originally published in 1981 under the name The Codex Standard, and updated in 2008 under the name The Codex Alimentarius. Each of the nutrients from breast milk and infant formula milk has advantages and disadvantages. The macronutrients and micronutrients contained in the nutrition provide support growth and development [15-17]. Late preterm babies take longer to reach the same size as full-term babies [18-20]. According to the research by Kakaroukas *et al.* (2022), feeding method was associated with fat mass, which can affect body weight and body length. Breastfed infant demonstrated greater increasing body weight than formula milk in late and moderate preterm infant [21]. In this study, we intended to look for differences in anthropometric status (body weight, body length, and head circumference) in late preterm babies. This work was performed in Dr. Soetomo General Hospital, Surabaya. It is the largest tertiary hospital in Eastern Indonesia. Sampling was carried out in March-June 2023.

## Experimental

### Material

Subjects were babies aged 6-12 months born at 34-37 weeks of gestation who came to the

pediatric inpatient and outpatient clinic at our hospital and had signed informed consent for research by the parents or caregivers. From a population of 106 babies, 70 samples were obtained. We excluded infants with birth weights  $\leq 1500$  and  $> 3000$  grams, any congenital abnormalities that affected measurements, large body masses, seriously ill infants, or infants with medical emergencies.

### Methods

This retrospective cohort study was conducted through interviews with parents or primary caregivers regarding age, gender, primary caregiver, and the main nutritional choices provided at the ages of 1, 3, and 6 months using a questionnaire created by the researcher. After that, we checked the anthropometric data obtained from each child's book. This work was approved by Komite Etik Penelitian Kesehatan, Dr. Soetomo General Hospital, Surabaya, Indonesia with the registered number 2050/105/3/III/2023.

We used IBM SPSS 24.0 statistics software. The normality test was carried out using Kolmogorov-Smirnov while the homogeneity test on normally distributed data used Levene. Statistical analysis was carried out using Kruskal-Wallis's test for the not normally-distributed data and the one-way Anova for normally and homogeneously distributed data.

## Results and discussion

### Demographic characteristics

This work is a retrospective cohort study that analyzes the difference in body weight, body length, and head circumference of late preterm babies fed with breast milk, infant formula milk, and a combination of breast milk and instant formula milk at Dr. Soetomo General Hospital Surabaya.

Late preterm babies are preterm babies born at 34-37 weeks of gestation [20]. The determination of gestational age is based on PMA (Post Menstrual Age) and ultrasound performed by professional health workers. The gold standard is first trimester ultrasound [22] In this study, 70 samples were obtained from a population of 106 late preterm babies who came to the pediatric outpatient and inpatient installation at dr. Soetomo General Hospital from March to June 2023. Babies born at 36-<37 weeks of gestation had the highest percentage (41.4%). Delnord *et al.* in 2019 who researched the epidemiology of late preterm births in several countries on the European, Asian, and American continents, stating that the prevalence of late preterm births aged 36-37 weeks in all countries studied most frequently among births at 34-35 and 35-36 weeks of gestation. In this study, the exact cause of the majority of late preterm births at 36-37 weeks was not definitively explained. [23] Saphiro-Mendoza in 2012, noted that multiple gestation, age of the mother (<20 and > 30 years), congenital malformation, and non-Hispanic black mother had experienced the most of late preterm birth in United States, it possibly caused by socio-economic issues. [22] However, the research indicated that a notable percentage of twin births occurred during this period (37.5%). A gestational age of 36-37 weeks is close to full-term gestational age. This can cause differences in the grouping of babies so that babies who should be in the term category become preterm. Several other studies include gender demographic data, but there is no discussion of gender influence on late preterm births. [Table 1]

Most of the subjects were raised by mothers (93.6%). Four subjects were raised by grandmothers and aunts, two subjects because the mother died, one subject because the mother was seriously ill, and one subject because the mother had a mental disorder. Parenting by mothers and other than parents is related to the maturity of psychosocial

development. Nurturing by mothers or biological parents influence children's maturity of psychosocial development. Maternal mood and psychological health can affect infant behavior and successful exclusive breastfeeding [24-27]. Handayani *et al.* in 2018, found that maternal factors were more important than infant factors to establish successful breastfeeding [28]. However, according to research conducted by Breheny *et al.* in 2013, children who are cared for by other than their parents will still have good psychosocial development if the caretaker implements good parenting [29].

The majority of mothers give breast milk to their newborn babies until one month of age (37 babies). The best nutrition for babies less than 6 months is exclusive breast milk [12,30,31]. Breast milk is considered the best nutrition for babies, especially late preterm babies because these babies are considered to have low levels of antioxidants in their bodies. Breast milk contains many immune components. In fact, according to ABM (Academy of Breastfeeding Medicine) recommendations, late preterm babies treated in the NICU should ideally be given breast milk from the mother or donor to optimize their growth and development. Breast milk is a source of bioactive protein, including enzyme activation, increased nutrient absorption, growth stimulation, immune system modulation, and defense against pathogens which is also known as antimicrobial activity [8,9,32-33]. In Indonesia, according to national economic survey in 2017, 55.96% babies fed with exclusive breast milk [34]. To improve exclusive breastfeeding and infant outcomes, mothers may need emotional support to understand how to breastfeed correctly, including techniques and how to obtain breast milk [24,35].

At birth, the most of babies were born with normal birth weight. The baby's weight normally decreases by as much as 10% of birth weight, then increases to normal weight.

Normal weight gain in preterm babies is 15 g/kgBW/day. Babies with normal birth weight, but at a lower limit, can become underweight babies. The early postnatal growth of late preterm infants is characterized by a major deposition of fat mass [12,18,37]. Martin *et al.* (2021), found that severe underweight babies are more common in premature babies than full term babies [37]. Differences in measurements and readings can also be the cause of changes in results. Gianni *et al.* (2012), late preterm babies can reach a weight at the corrected age of 3 months. However, according to Santos *et al.* (2009), late preterm babies tend to experience failure to thrive in the first 2 years of life [18,38].

The analysis of body weight from 1 month to 3 months of age could not be conducted due to the utilization of different growth charts.

The Fenton chart was employed at 0 and 1 months of age, whereas at 3 and 6 months of age, we referred to the 2006 World Health Organization (WHO) growth chart (Table 2).

There were 19 babies born with short stature (24.4%), while more babies were born with normal stature, 59 babies (75.6%). According to Brown *et al.* (2016) who studied 1071 premature babies, the average growth in length of babies was 0.12 cm/kgBW/day. At 3 months of age, the data was plotted using the 2006 WHO chart, resulting in 19 babies with severely stunted. There were 12 babies born with small head circumferences (15.4%). The normal growth of a baby's head circumference according to Brown *et al.* (2016) is 0.08 cm/week. According to Hallowell *et al.* (2012), an increase in head circumference is associated with the development of brain nerves/synapses [5,39].

**TABLE 1** Demographic characteristics of subjects

Variable	N (subjects) = 70	Percentage (%)
<b>Weeks of gestation</b>		
34-35	23	32.9
35-36	18	25.7
36-37	29	41.4
<b>Sex</b>		
Male	36	51.4
Female	34	48.6
<b>Caregiver</b>		
Mothers	66	94.3
Grand mother	4	5.7
<b>Nutrition at 1 month of age</b>		
Breastmilk	37	52.9
Infant Formula Milk	12	17.1
Combination of Both	21	30
<b>Nutrition at 3 month of age</b>		
Breastmilk	23	32.9
Infant Formula Milk	20	28.6
Combination of Both	27	38.6
<b>Nutrition at 6 month of age</b>		
Breastmilk	18	25.7
Infant Formula Milk	27	38.6
Combination of Both	25	35.7
<b>Birth Weight (BW)</b>		
Underweight	14	20
Normal	56	80
Overweight	0	0
<b>Birth Length</b>		
Severely Stunted	15	21.4
Normal	55	78.6
Tall	0	0

Variable	N (subjects) = 70	Percentage (%)
<b>Birth Head Circumference</b>		
Microcephaly	8	11.4
Normocephaly	62	88.6
Macrocephaly	0	0

**TABLE 2** Anthropometric Status late preterm baby in 0 and 1 month of age using Fenton growth chart

Variable	0 Month (Birth)		1 Month	
	n	%	n	%
<b>Body Weight</b>				
Underweight	14	20	49	70
Normal	56	80	21	30
Overweight	0	0	0	0
<b>Body Length</b>				
Short Stature	15	21.4	23	32.9
Normal Stature	55	78.6	47	67.1
Tall stature	0	0	0	0
<b>Head Circumference</b>				
Small	8	11.4	6	8.5
Normal	62	88.6	64	91.5
Big	0	0	0	0

The World Health Organization (WHO) 2006 growth chart is used for babies at 3 months and 6 months of age. At 3 months of age, 60% of subjects (42 babies) with normal weight, at 6 months of age the subjects with normal weight were increased to 60.3% of subjects (47 babies). At least 58.6% of subjects (41 babies) at 3 months of age were normal stature, but at the 6 months of age, the subjects with normal stature decreased to only 37.2% (26 babies). Based on HCAZ, 87.2% of subjects (61 babies) at 3 months of age were normal head circumference, but at 6 months of age, increased to 91.4% (64 babies) (Table 3).

*Differences in body weight of late preterm babies fed with breast milk, infant formula milk, and a combination of both*

One-way ANOVA statistical analysis showed that the body weight of late preterm babies aged 1,3, and 6 months at Dr. Soetomo General Hospital was not significantly different. These results mean that babies who were fed with breast milk, infant formula milk, and a combination of both had the same body

weight at 1, 3, and 6 months of age. At 3 months of age, it showed that babies who were given breast milk had the highest body weight compared to others, while babies who were given infant formula milk had the lowest body weight (Table 4).

At 6 months of age, babies who were fed with breast milk have the highest body weight compared to others, while babies who were fed with a combination of breast milk and instant formula milk had the lowest body weight. The subjects who were fed with breast milk, infant formula milk, or a combination of both had the same body weight at 6 months of age.

In this study, providing breast milk, infant formula milk, and a combination of both did not affect the weight of late preterm babies. Late preterm babies often appear to be full-term babies, but some babies can experience health problems. Stable or even decreasing body weight is one of the acute signs of nutritional problems in children. Preterm babies have accumulated problems with a lack of energy, proteins, minerals, and other nutrients when they first enter the NICU room. According to Gianni *et al.* (2016), late

preterm babies can lead to normal body weight in the first month of life because there is an accelerated increase in fat-free body mass composition. According to the Indonesian Pediatrician Association, preterm babies can experience weight gain in the normal body 15 g/kg/day. Meanwhile, according to ABM, the weight gain of late preterm babies can reach 20 g/day. According to research by Santos *et al.* in 2009, late preterm is one of the risk factors for growth failure and nutritional problems when the child is more than one year old [9,38,40-42]. This study is located in a tertiary hospital which is a referral hospital for babies experiencing health problems.

*Differences in body length of late preterm babies fed with breast milk, infant formula milk, and a combination of both*

At 1 month of age, the subjects who were fed breast milk had a higher average body length than babies who were fed with a combination of breast milk and instant formula milk, while babies who were fed with infant formula milk had the shortest body length. The results of the Kruskal-Wallis statistical analysis test showed that the body length at 1 month of age in our hospital was not significantly different. These results mean that babies who are fed with breast milk, infant formula milk, or a combination of breast milk and instant formula milk had the same body length at 1 month of age (Table 5).

At 3 months of age, statistical analysis test showed that the body length of the subjects 3 months of age in our hospital was not significantly different. These results mean that the subjects who were fed with breast milk, infant formula milk, or a combination of breast milk and instant formula milk had the same body length at 3 months of age

At 6 months of age, the subjects who were fed with breast milk had the highest average body length than the subjects who were fed with others, while babies who were fed with a

combination of breast milk and instant formula milk had the shortest body length. This shows that giving breast milk can increase the height of late preterm babies at 6 months of age. However, the results of the statistical analysis test showed that providing breast milk, infant formula milk, or a combination of breast milk and instant formula milk did not affect the body length of late preterm babies at 6 months of age.

Giving a combination of breast milk and instant formula milk to babies provides the shortest body length at 1 month, 3 months, and 6 months. Babies who were fed with breast milk had a higher body length than babies who were fed with a combination of breast milk and instant formula milk at the ages of 3 months and 6 months. When they reached 6 months of age, the subjects who were fed with infant formula milk had the highest body length, reaching 64.02 cm compared to babies who were fed with breast milk or a combination of breast milk and instant formula milk.

It was found that providing breast milk, infant formula milk, and a combination of both did not affect the body length of late preterm babies. However, according to Hallowell *et al.* (2016), preterm babies who experience mineral deficiency for a long time can cause metabolic bone disease and slow bone growth, so the baby becomes short [5]. Body length is an indicator of nutritional status better than body weight because it is not affected by the amount of fluid and reflects true lean body mass if done correctly. Preterm babies can experience an increase in body length of 0.8-1.0 cm/week. According to research conducted by Brown *et al.* (2016), late preterm babies who are fed breast milk with Human Milk Fortifier, can experience an increase in body length of 0.12 cm/week [39]. According to research by Santos *et al.* (2009), one of the risk factor for failure to thrive and nutritional problems when a child in more than one year old is late preterm birth [38]. Whereas, the first month after birth of late

preterm infants are characterized by rapid postnatal catch-up growth, therefore, at term-corrected age, they reach a weight that is either similar to or greater than that of term infant [40].

*Differences in the head circumference of late preterm babies fed with breast milk, infant formula milk, and a combination of both*

At 1 month of age, the babies who were fed with a combination of breast milk and instant formula milk had the largest head circumference compared to other nutrients, while babies who were fed with infant formula milk had the smallest head circumference. Kruskal-Wallis statistical analysis test showed that babies who were fed with breast milk, infant formula milk, or a combination of breast milk and instant formula milk had the same head circumference at 1 month of age (Table 6).

At 3 months of age, the subjects who were fed with a combination of breast milk and instant formula milk had the largest head circumference than others, while the subjects who were fed with infant formula milk had the smallest head circumference. The results of the statistical analysis test showed the babies who were fed with breast milk, infant formula milk, or a combination of breast milk and instant formula milk had the same head circumference at 3 months of age.

At 6 months of age, the subjects who were fed with infant formula milk had the largest head circumference than others, while the subjects who were fed with breast milk had

the smallest head circumference. Based on the results of Kruskal-Wallis statistical analysis test, the babies who were fed breast milk, infant formula milk, or a combination of breast milk and instant formula milk had the same head circumference at 6 months of age (Table 5).

It is showed that providing breast milk, infant formula milk, and a combination of both did not affect the head circumference of late preterm babies. An increase in head circumference indicates brain development. According to the Indonesian Pediatrician Association, the normal growth of head circumference in preterm babies is 0.5-0.8 cm/week. According to research by Brown *et al.* (2016), late preterm babies who are fed with exclusive breast milk with HMF can experience an increase in head circumference of 0.08cm/week compared to babies who are given exclusive breast milk without HMF [39]. According to research by Santos *et al.* in 2009, late preterm is one of the risk factors for failure to thrive and nutritional problems when the child is more than one year old [38]. Sammallahti in 2017, found that there is a correlation between growth spurt during the critical early period after birth (5-20 months) and better adult neurocognitive functioning. However, the same correlation is not consistently seen with mental health outcomes. This suggests that there may be different factors at play when it comes to the development of cognitive functioning versus mental health [43,44].

**TABLE 3** Anthropometric Status late preterm baby in 3 and 6 months of age using 2006 World Health Organization (WHO) growth chart

Variable	3 Months		6 Months	
	n	%	N	%
<b>WAZ</b>				
Severely Underweight	17	24.2	13	18.6
Underweight	11	15.7	14	20
Normal	42	60	43	61.4
<b>LAZ</b>				
Severely Stunted	15	21.4	30	42.8

Variable	3 Months		6 Months	
	n	%	N	%
Stunted	14	20	14	20
Normal	41	58.6	26	37.2
<b>HCAZ</b>				
Macrocephaly	1	1.4	0	0
Normal	61	87.2	64	91.4
Microcephaly	8	11.4	6	8.6

**TABLE 4** Body weight difference of late preterm baby in 1, 3, and 6 months of age

Birth Weight	n	Min-Max	Median	Mean	Std. Deviation	P-value
<b>1 Month<sup>(a)</sup></b>						
Breastmilk	37	1500-3900	2450	2578.81	580.98	0.795
Infant formula milk	12	1600-3900	2300	2538.46	698.26	
Combination of Both	21	1500-3800	2400	2508.70	617.85	
<b>3 Months<sup>(a)</sup></b>						
Breastmilk	27	2900-5700	4000	4024.07	821.28	0.365
Infant formula milk	22	2500-5700	3350	3581.82	947.99	
Combination of Both	29	2700-5000	3900	3748.28	651.71	
<b>6 Months<sup>(a)</sup></b>						
Breastmilk	20	4700-7800	6000	5935	787.58	0.868
Infant formula milk	30	3400-7800	5900	5756.67	1067.29	
Combination of Both	28	4300-7500	5350	5500	875.17	

(a) One-way ANOVA

**TABLE 5** Body length difference of late preterm baby in 1, 3, and 6 months of age

Variable	3 Months		6 Months	
	n	%	N	%
<b>WAZ</b>				
Severely Underweight	17	24.2	13	18.6
Underweight	11	15.7	14	20
Normal	42	60	43	61.4
<b>LAZ</b>				
Severely Stunted	15	21.4	30	42.8
Stunted	14	20	14	20
Normal	41	58.6	26	37.2
<b>HCAZ</b>				
Macrocephaly	1	1.4	0	0
Normal	61	87.2	64	91.4
Microcephaly	8	11.4	6	8.6

(k)Kruskal-Wallis

**TABLE 6** Head circumference difference of late preterm baby in 1, 3, and 6 months of age

Head Circumference	n	Min-Max	Median	Mean	Std. Deviation	P
<b>1 Month<sup>(k)</sup></b>						
Breastmilk	37	29-37	34	34.10	1.92	0.656
Infant formula milk	12	27-36	34	33.85	2.41	
Combination of both	21	32-38	34	34.61	1.62	
<b>3 Months<sup>(k)</sup></b>						
Breastmilk	23	32-41	37	37.19	2.10	0.656
Infant formula milk	20	28-40	37,5	37.16	2.45	



Head Circumference	n	Min-Max	Median	Mean	Std. Deviation	P
Combination of both 6 Months <sup>(k)</sup>	27	34-41	37	37.31	1.97	
Breastmilk	18	38-44	41	40.85	1.70	0.374
Infant formula milk	27	32-44	42	41.27	2.26	
Combination of both	25	35-46	41,5	40.89	2.42	

## Conclusion

This study aimed to analyze the difference in body weight and length. However, the results of this study were mostly not significant. Some of the weaknesses of this work following that the study was conducted in a tertiary hospital, that had a higher proportion of patients with medical problems both before and after birth. Moreover, this study used a retrospective cohort research method using secondary data source. The lack of standardized measuring instruments was observed. Measurements carried out by each health facility or Primary Health Care could be highly subjective during the execution and interpretation of results. In conclusion, there were no significant differences in weight, body length, and head circumference of late preterm babies fed with breast milk, infant formula milk, and combination of both aged 1,3, and 6 months. Other research about this study is needed, with standardized tools and method. Multicenter study may be preferred. It is hoped that the results of this research can be used as a consideration for providing nutrition to late preterm babies.

## Acknowledgments

We would like to appreciate the help of all staff in the Department of Child Health for supporting this study. We also like to thank the Head of Department of Child Health for approving our study.

## Funding

The authors declare no external funding.

## Authors' contributions

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved the final manuscript for publication.

## Conflict of interest

The authors declare no conflict of interests.

## Orcid:

Indana Lazulfa:

<https://orcid.org/0009-0001-4377-0844>

Martono Tri Utomo:

<https://orcid.org/0000-0003-4828-2134>

Nur Aisyah Widjaja:

<https://orcid.org/0000-0002-4253-8760>

## References

- [1] S. Asadi, FH Bloomfield, J.E. Harding, Nutrition in late preterm infants, *Seminars in Perinatology*, **2019**, *43*, 1–6. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [2] C.E. Briere, R. Lucas, J.M. McGrath, M. Lussier, E. Brownell, Establishing breastfeeding with the late preterm infant in the NICU, *Journal of Obstetric, Gynecologic & Neonatal Nursing*, **2015**, *44*, 102-113. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [3] P. Meier, A.L. Patel, K. Wright, J.L. Engstrom, Management of breastfeeding during and after the maternity hospitalization for late preterm infants, *Clinics in Perinatology*, **2013**, *40*, 689-705. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [4] X. Zhang, M. Zhou, H. Yin, Y. Dai, Y. Li, The predictive value of early oral motor assessments for neurodevelopmental outcomes of moderately and late preterm

- infants, *Medicine*, **2017**, 96, 9207. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [5] S.G. Hallowell, D.L. Spatz, The relationship of brain development and breastfeeding in the late-preterm infant, *Journal of Pediatric Nursing*, **2012**, 27, 154-162. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [6] J.B. Gouyon, S. Iacobelli, C. Ferdynus, F. Bonsante, Neonatal problems of late and moderate preterm infants, In *Seminars in Fetal and Neonatal Medicine* **2012**, 17, 146 -152. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [7] Z. Moudi, B. Molashahi, M. Imani, H. Ansari, Effects of a feasible supportive care program on breastfeeding behaviors and neonatal outcomes among the late preterm newborns in the south east of Iran, *Journal of Neonatal Nursing*, **2017**, 23, 238-241. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [8] B. Lönnerdal, Bioactive proteins in breast milk, *Journal of Paediatrics and Child Health*, **2013**, 49, 1-7. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [9] L.M. Noble, A.C. Okogbule-Wonodi, M.A. Young, A.O.B. Medicine, ABM clinical protocol# 12: transitioning the breastfeeding preterm infant from the neonatal intensive care unit to home, revised 2018, *Breastfeeding Medicine*, **2018**, 13, 230-236. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [10] S. Trend, T. Strunk, M.L. Lloyd, C.H. Kok, J. Metcalfe, D.T. Geddes, C. . Lai, P. Richmond, D.A. Doherty, K. Simmer, Levels of innate immune factors in preterm and term mothers' breast milk during the 1st month postpartum, *British Journal of Nutrition*, **2016**, 115, 1178-1193. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [11] W.G.W. Wardani, R.W. Winarni, Mendesain poster motivasi sebagai media edukasi anak-anak pemulung dan dhuafa di sekolah kami, bekasi barat, *RESWARA: Jurnal Pengabdian Kepada Masyarakat*, **2021**, 2, 365-378. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [12] M.L. Gianni, D. Consonni, N. Liotto, P. Roggero, L. Morlacchi, P. Piemontese, C. Menis, F. Mosca, Does human milk modulate body composition in late preterm infants at term-corrected age?, *Nutrients*, **2016**, 8, 664. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [13] J.V. Radtke, The paradox of breastfeeding-associated morbidity among late preterm infants, *Journal of Obstetric, Gynecologic & Neonatal Nursing*, **2011**, 40, 9-24. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [14] B.L. Crippa, L. Colombo, D. Morniroli, D. Consonni, M.E. Bettinelli, I. Spreafico, G. Vercesi, P. Sannino, P.A. Mauri, L. Zanotta, Do a few weeks matter? Late preterm infants and breastfeeding issues, *Nutrients*, **2019**, 11, 312. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [15] B. Koletzko, Z.A. Bhutta, W. Cai, S. Cruchet, M.E. Guindi, G.J. Fuchs, E.A. Goddard, J.B. Van Goudoever, S.H. Quak, B. Kulkarni, Compositional requirements of follow-up formula for use in infancy: recommendations of an international expert group coordinated by the Early Nutrition Academy, *Annals of Nutrition and Metabolism*, **2013**, 62, 44-54. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [16] M. Arendt, Codex Alimentarius: what has it to do with me?, *Journal of Human Lactation*, **2018**, 34, 704-710. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [17] A. Lapillonne, D.L. O'Connor, D. Wang, J. Rigo, Nutritional recommendations for the late-preterm infant and the preterm infant after hospital discharge, *The Journal of Pediatrics*, **2013**, 162, 90-100. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [18] M.L. Gianni, P. Roggero, N. Liotto, O. Amato, P. Piemontese, D. Morniroli, B. Bracco, F. Mosca, Postnatal catch-up fat after late preterm birth, *Pediatric Research*, **2012**, 72, 637-640. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [19] J.E. McGowan, F.A. Alderdice, V.A. Holmes, L. Johnston, Early childhood development of late-preterm infants: a systematic review, *Pediatrics*, **2011**, 127, 1111-1124. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [20] P. Gupta, R. Mital, B. Kumar, A. Yadav, M. Jain, A. Upadhyay, Physical growth, morbidity profile and mortality among healthy late

- preterm neonates, *Indian Pediatrics*, **2017**, *54*, 629-634. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [21] C.K. Shapiro-Mendoza, E.M. Lackritz, Epidemiology of late and moderate preterm birth. In *Seminars in Fetal and Neonatal Medicine*, **2012**, *17*, 120-125. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [22] Kakaroukas, M. Abrahamse-Berkeveld, L. Hayes, R.J. McNally, J.E. Berrington, R.M. van Elburg, N.D. Embleton, Early infancy growth, body composition and type of feeding in late and moderate preterms, *Pediatric Research*, **2023**, *93*, 1927-1935. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [23] M. Delnord, J. Zeitlin, February, Epidemiology of late preterm and early term births—An international perspective. In *Seminars in Fetal and Neonatal Medicine*, **2019**, *24*, 3-10. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [24] O.W. Edwards, School perceptions of children raised by grandparents, *Journal of Applied School Psychology*, **2018**, *34*, 86-100. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [25] S. Dib, J. C. Wells, M. Fewtrell, Mother And late Preterm Lactation Study (MAPLeS): a randomised controlled trial testing the use of a breastfeeding meditation by mothers of late preterm infants on maternal psychological state, breast milk composition and volume, and infant behaviour and growth, *Trials*, **2020**, *21*, 1-11. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [26] I. Estalella, J. San Millán, M. J. Trincado, A. Maquibar, L. Martínez-Indart, M. San Sebastián, Evaluation of an intervention supporting breastfeeding among late-preterm infants during in-hospital stay, *Women and Birth*, **2020**, *33*, 33-38. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [27] J.R. Demirci, S.M. Sereika, D. Bogen, Prevalence and predictors of early breastfeeding among late preterm mother–infant dyads, *Breastfeeding Medicine*, **2013**, *8*, 277-285. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [28] K.D. Handayani, Irwanto, M. Masturina, R. Etika, A. Harianto, P.J. Sauer, Duration of breastfeeding in late preterm infants: maternal and infant factors, *Journal of Human Lactation*, **2021**, *37*, 795-802. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [29] M. Breheny, C. Stephens, L. Spilsbury, Involvement without interference: How grandparents negotiate intergenerational expectations in relationships with grandchildren, *Journal of Family Studies*, **2013**, *19*, 174-184. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [30] A. Keir, A. Rumbold, C.T. Collins, A.J. McPhee, J. Varghese, S. Morris, T.R. Sullivan, S. Leemaqz, P. Middleton, M. Makrides, Breastfeeding outcomes in late preterm infants: a multi-centre prospective cohort study, *Plos One*, **2022**, *17*, 272583. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [31] E. Post, G. Stam, E. Tromp, Milk production after preterm, late preterm and term delivery; effects of different breast pump suction patterns, *Journal of Perinatology*, **2016**, *36*, 47-51. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [32] M. Forsgren, E. Isolauri, S. Salminen ,S. Rautava, Late preterm birth has direct and indirect effects on infant gut microbiota development during the first six months of life, *Acta Paediatrica*, **2017**, *106*, 1103-1109. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [33] A.I. Eidelman, Breastfeeding and the use of human milk: an analysis of the American Academy of Pediatrics 2012 Breastfeeding Policy Statement, *Breastfeeding Medicine*, **2012**, *7*, 323. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [34] J.Y. Meek, L. Noble, Breastfeeding and the use of human milk, *Pediatrics*, **2022**, *150*, 2022057988. [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [35] S.Y. Lee, G.J. Jang, S.Y. Lee, G.J. Jang, Prevalence and predictors of exclusive breastfeeding in late preterm infants at 12 weeks, *Child Health Nursing Research*, **2016**,

- 22, 79-86. [Crossref], [Google Scholar], [Publisher]
- [36] R.M. Kliegman, B.F. Stanton, J. St Geme, N.F. Schor, *Nelson Textbook of Pediatrics*; Elsevier Health Science: Amsterdam, 2015; p. 1128-1132.
- [37] M. Gómez-Martín, D. Herrero-Morín, G. Solís, M. Suarez, N. Fernández, S. Arboleya, M. Gueimonde, S. González, Longitudinal study depicting differences in complementary feeding and anthropometric parameters in late preterm infants up to 2 years of age, *Nutrients*, **2021**, *13*, 982. [Crossref], [Google Scholar], [Publisher]
- [38] I.S. Santos, A. Matijasevich, M.R. Domingues, A.J. Barros, C.G. Victora, F.C. Barros, Late preterm birth is a risk factor for growth faltering in early childhood: a cohort study, *BMC Pediatrics*, **2009**, *9*, 1-8. [Crossref], [Google Scholar], [Publisher]
- [39] J.V.E. Brown, L. Lin, N.D. Embleton, J.E. Harding, W. McGuire, Multi-nutrient fortification of human milk for preterm infants, *Cochrane Database of Systematic Reviews*, **2020**, [Crossref], [Google Scholar], [Publisher]
- [40] M.L. Gianni, E. Bezze, P. Sannino, E. Stori, L. Plevani, P. Roggero, M. Agosti, F. Mosca, Facilitators and barriers of breastfeeding late preterm infants according to mothers' experiences, *BMC Pediatrics*, **2016**, *16*, 1-8. [Crossref], [Google Scholar], [Publisher]
- [41] A. Protocol, ABM Clinical Protocol# 10: Breastfeeding the Late Preterm (34–36 6/7 Weeks of Gestation) and Early Term Infants (37–38 6/7 Weeks of Gestation), Second Revision 2016, *Breastfeeding Medicine*, **2016**, *11*, [Crossref], [Google Scholar], [Publisher]
- [42] K.I.D.A. Indonesia, Asuhan Nutrisi pada Bayi Prematur, *Konsensus*, **2016**, *5*. [Google Scholar], [Publisher]
- [43] S. Sammallahti, K. Heinonen, S. Andersson, M. Lahti, S. Pirkola, J. Lahti, A.-K. Pesonen, A. Lano, D. Wolke, J.G. Eriksson, Growth after late-preterm birth and adult cognitive, academic, and mental health outcomes, *Pediatric Research*, **2017**, *81*, 767-774. [Crossref], [Google Scholar], [Publisher]
- [44] M. Woythaler, February. Neurodevelopmental outcomes of the late preterm infant, In *Seminars In Fetal and Neonatal Medicine*, **2019**, *24*, 54-59. [Crossref], [Google Scholar], [Publisher]

**How to cite this article:** Indana Lazulfa, Martono Tri Utomo, Nur Aisiyah Widjaja, Differences in anthropometric status in late preterm babies fed with breastmilk, infant milk, and a combination of both. *Journal of Medicinal and Pharmaceutical Chemistry Research*, 2024, 6(11), 1767-1778. **Link:** [https://jmpcr.samipubco.com/article\\_196694.html](https://jmpcr.samipubco.com/article_196694.html)